



Global Advanced Research Journal of Agricultural Science (ISSN: 2315-5094) Vol. 4(2) pp. 089-094, February, 2015.  
Available online <http://garj.org/garjas/index.htm>  
Copyright © 2015 Global Advanced Research Journals

*Full Length Research Paper*

# Effect of three covering materials on vegetative growth of cucurbits in Kenya.

**Arnold Onyango Watako**

Department of Agricultural Economics and Agribusiness Management, School of Agricultural and Food Sciences; Jaramogi Oginga Odinga University of Science and Technology, P.O Box 210 – 40601, Bondo, Kenya.  
Email: [arnoldwatako@yahoo.com](mailto:arnoldwatako@yahoo.com)

Accepted 15 January, 2015

Different covering materials were compared at Jomo Kenyatta University of Agriculture and Technology, Juja in Kenya ( $1^{\circ} 5' S$  latitude,  $37^{\circ} 1' E$  longitude and 1520m above sea level) on their effect on vegetative growth of cucurbits. Three structures equivalent of main treatments each having ventilation gaps were covered with 50% shade net ( $T_1$ ); glass ( $T_2$ ) and UV stabilized plastic sheet ( $T_3$ ) materials respectively were used. The sub treatments were cucurbits family; cucumber and butternut respectively. In each structure twelve pots were placed, six of each crop type. The experiment was laid out as a random complete block design and replicated three times, each replication consisting of two pots. Dry bulb temperature was highly significant in all the treatments with plastic house having the highest mean temperature of  $33.83^{\circ}C$  followed by glass house with  $32.5^{\circ}C$  and shade house with  $30^{\circ}C$  respectively while the wet bulb temperature was not significantly different at  $p < 0.05$ . The relative humidity in all the treatments was not significantly different at  $p < 0.05$  although plastic house had the highest RH of 91.5%, followed by shade house with 90.5% and glass house with 80%. There were significant differences in weeks 3, 6, 7 and 8 for plant height but no significant influence with the type of covering material. However, shade house treatment dominated cumulatively in plant height. Glasshouse treatment yielded most leaves in week 4 for cucumber, weeks 6 and 7 for butternut and week 8 for both butternut and cucumber. Solar rays were more direct to the plant in shade house than in glasshouse or plastic house. There was a positive correlation between light intensity and leaf area ( $r=0.1$ ) showing that for every unit increase in light intensity, the leaf area increased by 0.1. Considering the temperature and relative humidity variables and the direct effect of solar rays in the shade house coupled with plant height parameter, shade house is an appropriate structure for cucurbit production in a tropical climate.

**Keywords:** Protected cultivation, cucurbitaceae, growth, Kenya

## INTRODUCTION

In Kenya, the vegetable sub – sector is important in attaining food security and improving livelihood for smallholder farmers. Smallholders produce 100% of African Leafy Vegetables (ALVs) and up to 70% of the Exotic and Asian vegetables (HCDA, 2013). The production of vegetable crops under structures covered with materials commonly referred to as protected

cultivation is gaining popularity in Kenya. Protected cultivation is the production of crops under sheltered structures covered with such materials as glass, plastic films or shade screens (saran cloth) which create a specialized climate inside the structure. Protected cultivation of vegetable crops suitable for domestic and export purposes could be a more efficient alternative for

land use and other resources, improve yield, quantity and quality (Sanwal et al., 2004; Ganesan, 2004). However, the profitability in protected cultivation depends upon the choice of structures, selection of crops/ varieties, production technology and market price.

Changes in the microclimate inside the structures result in modification of plant growth and development and also incidence of fungal diseases (Beckmann et al., 2006; Scaranari et al., 2008; Chavarria et al., 2009). In one study, interactions between protected environments and different cucumber hybrids were evaluated by Costa et al., (2010) and found that the seedling growth and dry biomass was affected by the growing environments. In greenhouse tomato production, Gent (2007) found that the use of the screen with 50% shading increased commercial fruit production by 9% compared to the environment without the screen. Comparisons between aluminized shade screens with 40, 50 and 60% shading and polyethylene plastic film painted with lime on tomato production showed that the screens minimize energy consumption during periods of low temperatures (Fernandez-Rodriguez et al., 2001). In acclimatized environment with evaporative cooling, Costa and Leal (2008) found greater accumulation of leaf biomass and greater leaf area of strawberry plants than in non acclimatized environments, regardless of the season. Shading with 40% shade net was beneficial for lettuce production (Queiroz et al., 2009).

Cucumber and butternut are members of the cucurbitaceae family. The potential for increased production is immense for these cucurbit crops in the on – going rehabilitation and expansion of irrigation schemes in arid and semi – arid counties in Kenya. In the present scenario of perpetual demand for vegetables and drastically shrinking land holdings, protected cultivation is the best alternative for using land and other resources more efficiently. There is an urgent need to assess the suitability of different covering materials for cultivation of vegetables to meet the growing demand. This study investigated the effect of shade net, plastic and glass covering materials on vegetative growth of cucurbits.

## MATERIALS AND METHODS

Different covering materials were compared for their effect on vegetative growth of cucurbits at Juja, Kenya during the periods between 26<sup>th</sup> June 2013 and 14<sup>th</sup> May 2014. The farm is located at 1° 5' S latitude, 37° 1' E longitude and 1520m above sea level. Three structures each having ventilation gaps were covered with 50% shade net; glass film and UV stabilized plastic sheet materials respectively. The soil media used was a cocktail of garden soil, manure and sand mixed in the ratio of 3:2:1 and put into poly tubes measuring 19cm in diameter and 21cm height. Five (5) gms of Diammonium phosphate (DAP) was placed in the soil media per pot, thoroughly mixed then watered. Seeds

of butternut pumpkin (*Curcubita moschata*) variety 'squash' and cucumber (*Cucumis sativus*) variety 'Ashley' were sown in thirty six pots; eighteen pots of each crop. The poly tubes were then left in the open field for the seeds to germinate which took approximately seven days before being transferred to the protective structures. Each structure contained six pots of butternut and six of cucumber giving a total of twelve pots. The treatments were thus;

T<sub>1</sub> – Shade house which was fully covered with 50% shade net with the door kept open for ventilation during day time

T<sub>2</sub>- The glasshouse consisted of the entire roof and four sides covered with glass but the sides alternated with glass and wire mesh.

T<sub>3</sub> – The plastic house was made up of the four sides and the roof covered with polyethylene film with side ventilation

In each structure, the experiment was laid out as a randomized complete block design and replicated three times. Each replication consisted of two pots. Standard cultural practices for cucurbit production were followed.

## Data taking and Analysis

Data for solar radiation was taken using a luximeter while humidity and temperature data were taken using a wet and dry thermometer (Figure 1 and 2). The side for measuring temperature was kept wet everyday by adding water in the container with the wick. Plant height was taken using a ruler and physical counting of leaves was done to determine the number of leaves per plant. Leaf size was determined by measuring the leaf lamina horizontally and vertically thus the leaf area was obtained by multiplying the horizontal and then vertical measurements of the leaf lamina. Horizontal measurements entailed measuring the widest parts of the leaf lamina and vertical measurements were obtained by measuring from the tip of the leaf to where the leaf stalk joined the lamina. Data collected was entered into Microsoft spreadsheets and subjected to computer statistical analysis using Genstat VSN International version 14.

## RESULTS

### Temperature

Dry bulb temperature was highly significant in all the treatments with plastic house having the highest mean temperature of 33.83°C followed by glass house with 32.5°C and Shade house with 30°C respectively at p<0.05 (Table 1). Similarly, the wet bulb temperature was significantly different for all the treatments. Nevertheless, plastic house still had the highest mean temperature of 32.5°C followed



Figure 1. Luximeter



Figure 2. Wet and dry bulb thermometer

Table 1. Means of the Wet and dry bulb temperature ( $^{\circ}\text{C}$ ), Luxmeter (Lux) and relative humidity (RH) for the three treatments

Treatment	Wet $^{\circ}\text{C}$	Dry $^{\circ}\text{C}$	RH(%)	Lux
Shade house	26c	30b	90.5a	440a
Glass house	29.33b	32.5c	80a	306b
Plastic house	32.5a	33.83a	91.5a	251c
Grand mean	29.28	32.11	87.2	333
CV%	11.5	3.6	15.2	19.4
L.S.D	4.13	1.4	16.32	79.2

Means with the same letters are not significant at  $p < 0.05$

by glass house with  $29.33^{\circ}\text{C}$  and Shade house with  $26^{\circ}\text{C}$ , respectively.

### Light intensity

The luxmeter readings in all the treatments was significantly different at  $p < 0.05$  with the highest light intensity being in shade house (440 lux) and the lowest was in plastic house (251 lux) (Table 1).

### Relative humidity

The relative humidity in all the treatments was not significantly different at  $p < 0.05$  although plastic house had the highest relative humidity of 91.5%, followed by shade

house with 90.5% and glass house with 80% (Table 1). In the glasshouse treatment there were cases of blights on leaves and the fruit that formed aborted due to fungal infection (Figure. 3).

### Plant height

There were significant differences within varieties in weeks 3, 6, 7 and 8 but there were no significant differences with the type of covering material. Interaction between Variety and treatment was not significant in all the weeks (Figure. 4). Butternut dominated in height in shade house treatment in week 3 and glasshouse in weeks 6, 7 and 8. Cucumber was the tallest in week 4 only although shade house



Figure 3. Butternut leaf affected with blight




Figure 4. The influence of covering material on plant height increase from week 3 to week 8  
Means with the same letters are not significantly different at  $p < 0.05$

treatment dominated in all the weeks with 2.83, 7, 10.67, 13.7, 28 and 51cm, respectively.

### Number of leaves

There were significant differences within the varieties of cucurbits in all the treatments as the number of weeks increased at  $P < 0.05$  (Figure 5). But there were no significant differences within the treatments at week level. Butternut had the highest number of leaves from week 3 to week 7 in all the treatments while cucumber had the highest number of leaves in all the treatments in week 8.

Glasshouse treatment yielded most leaves in week 4 for cucumber; weeks 6 and 7 for butternut and week 8 for both butternut and cucumber (Figure 5).

### Leaf area

There was a positive correlation between light intensity and leaf area ( $r = 0.1$ ) showing that for every unit increase in light intensity, the leaf area increased by 0.1 (Figure 6). Increase in leaf area directly affected plant growth and hence plant height.




Figure 5. The number of leaves per treatment from week 3 to week 8

Key: B- Butternut  
C-Cucumber

## DISCUSSION

Among environmental factors, light intensity, temperature and relative humidity influence crop growth and development in protected cultivation (Rajasekar et. al. 2013). The influence of environmental variables; temperature, relative humidity and light intensity under different covering materials were studied. The temperature in the plastic house was the highest probably due to the thermal transmissivity characteristics of plastic film. Despite plastic house treatment having significant temperatures above shade house and glasshouse respectively, dry matter accumulation that could have resulted into significant plant height was not achieved. This could be explained probably due to the fact that high temperatures which could have slowed down the cell metabolic activities thus lowering the growth rate. Similar studies with radish (*Raphanus sativus* L.) showed that higher temperatures had more adverse influence on net photosynthesis than lower temperatures leading to decreased production of photosynthates (Reddy et. al. 1999). Solar rays were more direct to the plants in shade house than in glass house or plastic house. In plastic house the light was more diffused and opaque as compared to the other two treatments. Despite the Shade house obtaining on average higher light intensity than other treatments (glasshouse and plastic house), the growth rate determined by plant height in this case was not significant.

A high relative humidity on the one hand encourages fungal diseases (Figure 3), because under fluctuating temperatures and sharply increasing evaporation during

the first morning hours, condensation can easily occur on the crop creating the ideal conditions for fungal spores to germinate rapidly. A high relative humidity can also cause a crop to weaken and become more susceptible to changes in the weather. High relative humidity could have contributed to slower growth rate because the air was almost at the saturation point and thus the respiration rate was slowed down. If the respiration rate is slowed down, there's accumulation of photosynthates within the tissues thus this also leads to lower photosynthetic rate and hence slower growth rate. This explains the reason why there were lower number of leaves in this treatment as compared with glasshouse and shade house (Figure 5) and also lower leaf area despite the high light intensity (Figure 6). Cucumber did not exhibit significant differences in the treatments applied. Shade house treatment dominated cumulatively in plant height because of its optimal temperature of 26°C, high relative humidity of 90.5% and high light intensity that supported growth and development. This observation supports the findings of Marcelis and Baan Hofman – Eijer (1993) which indicated that cucumber require more light intensity and high temperatures for better growth and development. A larger leaf area signified a larger proportion for light interception and since light is the critical energy for photosynthesis, the more the light translates to more photosynthates. These results agree with Papadopoulos and Omrod (1991) that higher leaf area increases leaf physiology and number of stomata and thus photosynthesis. This therefore leads to faster and higher growth rate as is the case in glasshouse (Figure 3). The higher the photosynthetic rate, the faster the phenological

events taking place in a plant thus also the better the quality of the plant. Therefore this also could have caused the glasshouse treatment which had a lux of 306 (Table 1) to have the highest number of leaves (Figure 5).

Ultimately, this study revealed that the prospect of cucurbits cultivation under shade house is promising. Other structures could also be used but only with sufficient ventilation to regulate the temperature and the humidity. In conclusion, the best protective covering for production of cucurbits and generally vegetables in areas of high temperatures and light intensity could be the Shade net. However, varietal interaction with the type of cover used needs investigation.

#### REFERENCES

- Beckmann MZ, Duarte GRB, Paula VA, Mendez MEG, Peil RMN (2006). "Radiação solar em ambiente protegido cultivado com tomateiro nas estações verão-outono do Rio Grande do Sul." *Ciência Rural*, Santa Maria-RS, v. 36, n. 1, p. 86-92, ISSN 0103-8478
- Chavarría G, Santos HP, Mandelli F, Marodin GAB, Bergamaschi H, Cardoso LS (2009). "Potencial produtivo de videiras cultivadas sob cobertura de plástico".
- Costa E, Leal PAM (2008). "Avaliação de alface hidroponica em três ambientes de cultivo". *Engenharia Agrícola*, Jaboticabal, v. 29, n. 3, p. 358 – 369, ISSN 0100 – 6916.
- Costa E, Leal PAM, Gomes VA, Machado D, Jara MC (2010). Biomassa de mudas de pepines híbridos conduzidos sob ambientes protegidos. *Bregantia*, Campinas, v. 69. N. 2, p. 381 – 386, ISSN 0006 – 8705.
- Fernandez – Rodriguez EJ, Perez D, Camacho- Ferre F, Fernandez VJ, Kenig A (2001). "Effects of aluminized shading screens vs whitewash on tomato photochemical efficiency under a non heated greenhouse". *Acta Hort.* v. 559, p. 279 – 284. ISSN 0567 – 7572.
- Ganesan M (2004). "Effect of poly – greenhouse on plant micro – climate and fruit yield of sweet pepper". *International Symposium on Integrated Management Practice*. AVRDC. Taiwan.
- Gent MPN (2007). "Effect of shade on quality of greenhouse tomato". *Acta Hort.* v. 747, p. 107 – 112, ISSN 0567 – 7572.
- HCDA (2013). "Horticulture validated report". Horticultural Crops Development Authority, Kenya; [www.hcda.or.ke](http://www.hcda.or.ke)
- Marcelis LFM, Baan Hofman – Eijer LR (1993). "Effect of temperature on growth of individual cucumber fruits". *Physio. Plantarum* 87: 321 – 328.
- Papadopoulos AP, Omrod DP (1991). "Plant spacing effects on growth and development of greenhouse tomato". *Canadian J. Plant Sci.* 71: 297 – 304.
- Pesquisa agropecuária brasileira, Brasília, v. 44, n. 2, p. 141-147, ISSN 0100-204X.
- Queiroz JPS, Neves LG, Seabra Junior S, Costa AJM (2009). "Avaliação da produção de genótipos de alface em diferentes ambientes, cultivados no período de inverno em Caceres/ MT." In *Jornada Científica da Unemat, Barra dos Burges - MT. Resúmenes.. Barra dos Burges – MT; Universidade Estadual de Mato Grosso*, 2009.
- Rajasekar M, Arumugam T, Rramash K (2013). "Influence of weather and growing environment on vegetable growth and yield." *Journal of Horticulture and Forestry*. Vol. 5(10); pp. 160 – 167.
- Reddy MT, Ismail S, Reddy YN (1999). "Shade and allelopathic effects of ber on growth, productivity and quality of radish ( *Raphanus sativus* L.) under pot culture." *South Indian Hortic.* 47: 77 – 80.
- Sanwal SK, Patel KK, Yadav DS (2004). "Vegetable production under protected conditions in NEH region." *Problems and Prospects. Indian Soc. Veg. Sci.* 3: 120 – 129.
- Scaranari C, Leal PAM, Pellegrino GQ (2008). "Estudo de simulações de microclimas em casas de vegetação visando à aclimação de mudas micropropagadas de bananeira cv Grande Naine." *Revista Brasileira de Fruticultura, Jaboticabal-SP*, v. 30, n. 4, p. 1001-1008, ISSN 0100-2945