

Global Advanced Research Journal of Food Science and Technology (ISSN: 2315-5098) Vol. 3(5) pp. 141-154, October 2014
Available online <http://garj.org/garjfst/index.htm>
Copyright © 2014 Global Advanced Research Journals

Full Length Research Paper

Effects of different maize (*Zea mays* L.) – soybean (*Glycine max* (L.) Merrill) intercropping patterns on yields, light interception and leaf area index in two contrasting sites

Jossias Mateus Materusse Matusso^{1*}, Jayne Njeri Mugwe², Monicah Mucheru-Muna³

^{1*}International Crops Research Institute for the Semi-Arid Tropic, P. O. Box 1906, Maputo, Mozambique.

²Department of Agricultural Resource Management, School of Agriculture and Enterprise Development, Kenyatta University (KU), Nairobi, P. O. BOX 43844 – 00100, Kenya

³Department of Environmental Science, School of Environmental Studies, Kenyatta University (KU), Nairobi, P. O. BOX 43844 – 00100, Kenya

Accepted 22 March 2014

Field trials were conducted at the field units of the Embu Agricultural Training Center and Kamujine Dispensary in Embu and Meru Counties, Kenya, during 2012 long rain (LR) and short rain (SR) seasons to determine the effects of different maize-soybean intercropping patterns on yields, light interception and leaf area index. The main treatments were four maize – soybean intercropping patterns (conventional-1maize:1soya; MBILI-2maize:2soya; 2maize:4soya; 2maize:6soya) and two sole crops of maize and soybean, respectively. The experimental design was a randomized complete block design with four replications, and plot size of 7.0 m by 4.5 m. The study was carried out in two seasons (long rain 2012 and short rain 2012). The results showed that, the maize-soybean intercropping patterns had significant effect on maize stover and grain yields during both seasons and sites. The MBILI treatment recorded significantly higher stover and grain yields than all other treatments. During the long rain 2012, the soybean yields were reduced by 60 and 81% due to the intercropping with maize, at Embu and Kamujine, respectively; whereas during the 2012 SR, the yields were reduced by 52 and 78% as effect of intercropping with maize at Embu and Kamujine sites, respectively. The intercropping patterns affected significantly ($p < 0.0001$) the photosynthetically active radiation intercepted and the leaf area index at both sites. From the results of this study, the use of MBILI maize-soybean intercropping pattern can be recommended to the farmers of central highlands of Kenya because it gave efficient resources use and higher yields.

Keywords: Intercropping patterns, maize-soybean, leaf area index (LAI), photosynthetically active radiation (PAR), central highlands, Kenya.

INTRODUCTION

In intercropping system there is one main crop cultivated

with one or more added crops where the main crop is of primary importance due to economic or food production reasons (Brintha and Seran, 2009). In the SSA region, cereal and grain legumes intercrop is the most practiced

*Corresponding Author Email: matujossias@gmail.com

by smallholder farmers (Odendo, Bationo and Kimani, 2011; Sanginga and Woomeer, 2009). The major reason why these farmers intercrop cereals and grain legumes is because they are particularly important human food as they are rich in protein and are sometimes sold for cash income (Odendo, Bationo and Kimani, 2011). In addition, intercrops give them the stability of the yields over several seasons (Ofori and Stern, 1987; Steiner, 1982), when one crop fails, the other might still give a reasonable yield (Prasad and Brook, 2005; Beets, 1982; Steiner, 1982). Furthermore, grain legumes help maintain and improve soil fertility due to their ability to biologically fix atmospheric nitrogen (Sanginga and Woomeer, 2009; Jarenyama *et al.*, 2000). Despite that, the intercropping of cereal-legume may lead to reduction in yield of the legume component because of the adverse competitive effects (Willey *et al.*, 1983). Often, the cereal component with relatively higher growth rate, height advantage and a more extensive rooting system is favored in the competition with the associated legume crop. Thus, the greater yield loss of the minor crop is mainly due to reduced photosynthetically active radiation (PAR) reaching the lower parts of the intercrop canopy, occupied by the minor legume (Liu *et al.*, 2010). The intensity and the quality of solar radiation intercepted by the canopy are important determinants of yield components and therefore yield of soybean since it is sensitive to shading (Liu *et al.*, 2010; Purcell, 2000). Light levels during the late flowering to mid pod formation stages of growth have been found to be more critical than during vegetative and late reproductive periods (Liu *et al.*, 2010; Schou, Jeffers and Streeter, 1978). Therefore, any interventions that lead to increased amount of PAR interception by the minor crop have potential to increase the yield of the minor crop and increase productivity of the intercropping system (Mashingaidze, 2004). For instance, Woomeer and Tungani (2003) reported that MBILI system had resulted in 20 percent more light penetration to the minor legume component when compared to the conventional intercropping pattern. Ennin, Clegg and Francis (2002) found that per cent PAR intercepted by intercrops was 4 percent greater in closer row arrangements of soybean and maize than in equally spaced 2 rows soybean: 2 rows maize. Reddy, Floyd and Willey (1980) reported that millet-groundnut intercrop system was 28 percent more efficient in light use than their monocrops, which was attributed to approximately 30 percent greater LAI of the intercrop than the sole crops. Leaf area index of a canopy is important for predicting crop growth and yields (Xinyou *et al.*, 2003). A reasonable LAI is critical to maintain high photosynthetic rates and the yield (Xiaolei and Zhifeng, 2002). If the index is too low, not enough light will be absorbed and if too high, lower leaves will not receive enough light and will thus be a liability (Brintha and Seran, 2009). On the other hand, intercropping can lead to reduction in yield of one or more of component crops due to adverse

competitive effects (Willey and Rao, 1980). A review by Ofori and Stern (1987), of 40 published papers showed that the yield of the legume component declined on average by about 52 percent of the sole crop yield whereas the cereal yield was reduced by only 11 percent. The general observations from this are that yields of the legume components are significantly depressed by cereal components in intercropping, which is attributed to reduced photosynthetically active radiation (PAR) that reaches the lower parts of the maize canopy occupied by the soybean crop.

We investigated the effects of different maize-soybean intercropping patterns on yields, light interception and leaf area index. The hypotheses tested were: (i) the MBILI intercropping pattern gives significantly higher yields than conventional intercropping pattern. (ii) the 2:6 intercropping pattern of maize-soybean is more efficient in light use and gives higher leaf area index than conventional (1M:1S) intercropping pattern.

MATERIALS AND METHODS

Study area

The experiment was carried out in two sub counties of central highlands of Kenya, namely Embu West and Tigania East sub counties.

Embu West Sub County

Embu West District is located in Embu County, in the central highlands of Kenya, and occupies an area of 708 Km² and is bordered by Mbeere district to the East and South East, Kirinyaga to the West and Meru South to the North. The experimental site lies within N 0° 31' 4.2'' E 37° 27' 20'' and at the altitude of 1468 m above the sea level (ASL), at Embu Agricultural Staff Training College. The major agro-ecological zone (AEZ) is Upper Midland 2 (UM 2). The soils are mainly humic Nitisols and the total arable land area is 478 Km² with total available agricultural land area covering 371 Km². The average annual rainfall varies from 909 to 1230 mm with long rainy season between March and June and short rainy season between October and December, respectively (Jaetzold *et al.*, 2006).

Tigania East Sub County

Tigania East Sub County is located in Meru County, in the central highlands of Kenya and it occupies 108.6 km². The experimental site lies within N 0° 6' 19.5'' E 037° 64' 39.6'' and at the altitude of 935 m above the sea level (ASL), at Kamujine Dispensary in Mikinduri Division. The major agro-ecological zones are Lower Midlands 3 and

Upper Midland 3 (LM 3 and UM 3), the soils are mainly eutric Nitisols and humic Cambisols. The annual average temperature varies from 19.2 °C to 22.9 °C. The average annual rainfall varies from 1000 to 2200 mm with long rainy season between March and June and short rainy season between October and December, respectively (Jaetzold *et al.*, 2006).

Experiment design and treatments

The experiment in this study was established in Embu-ATC (Embu West district) and in Kamujine (Tigania East district) in March 2012 and it was laid out as a randomized complete block design (RCBD). There were four replicate blocks and plot sizes measuring 7 m x 4.5 m. The cropping system was of sole maize (*Zea mays* L.), sole soybean (*Glycine max* (L.) Merrill) and maize (M) – soybean (S) intercropping with cropping patterns (Table 1).

Management of the experiment

The fields were ploughed using hand hoe and left as such for two weeks. Plots measuring 7.0 x 4.5 m were marked just before planting. Pathways measuring 3.0 m and 2.0 m were left between the blocks and plots, respectively. At Embu-ATC, planting was done on the 23rd of March and 12th of October 2012 for the 1st and 2nd seasons, respectively. At Kamujine, planting was done on the 26th of March and 15th of October 2012 for the 1st and 2nd seasons, respectively. The sole maize (*Zea mays* L.) var. DK 8031 was planted at a spacing of 0.75 m x 0.50 m inter and intra-row, respectively. The number of hills per row was 10 with three seeds per hill in order to ensure maximum plant population and to account for germination failure; and two weeks after germination the excess plants were thinned out to remain with two plants per hill. The sole soybean (*Glycine max* (L.) Merrill) var. Gazelle was hand drilled at a spacing of 0.45 m x 0.10 m in inter and intra – row spacing resulting to 62 plants per row to ensure maximum germination/population and the excess plants were thinned out to remain with the recommended population of 31 plants per row after 2 weeks of emergence. The following external nutrient replenishment inputs were applied per plot: 6kg of manure equivalent to 30 kg N ha⁻¹, applied two weeks before planting; 94.5 grams of CAN as source of N, equivalent to 30 kg N ha⁻¹, for soybean the Nitrogen (starter N) was applied at sowing while for maize it was applied when the crop had six leaves, as topdressing; 189 grams of TSP as source of P, equivalent to 60 kg P ha⁻¹, which was applied at sowing. The fertilizers were applied accordingly to the recommendation. Management practices were the same for both the monocrop and the maize – soybean intercrop.

Maize and soybean harvest and yields

Maize and soybean grain and stover was harvested at maturity from a net area of each treatment demarcated after leaving out two rows on each side of the plot and the first two and the last two maize/soybean plants on each row to minimize the edge effect. The entire plants on the plots was harvested by cutting at the ground level and weighted to represent the total fresh weight. Maize/Soybean cobs/pods were manually separated from the stover, sun-dried, and packed in sacks before threshing. After threshing, moisture content of the grains was determined using a moisture meter and grain yield adjusted to 12 percent moisture content using the following formula. Similarly, the yields were calculated using the following formulas.

$$\text{Adjusted yield} = \text{measured yield} * \frac{(100 - \text{sample moisture content})}{(100 - \text{standard moisture content})} \quad (1)$$

$$\text{Yield (t/ha)} = 10 * \frac{\text{Dry weight (kg/m}^2\text{)}}{\text{Net area (m}^2\text{)}} \quad (2)$$

Determination of light interception and leaf area index

Radiation interception of photosynthetically active radiation was measured in both the sole crop and the different intercropping patterns using a Sunfleck Ceptometer. The measurements were taken between 11:30 am and 01:30 pm (local time) at an interval of fourteen days. The PAR intercepted was then calculated according to Goudriaan (1988) as follows;

$$\% \text{ PAR intercepted} = \frac{(PAR_a - PAR_b)}{PAR_a} * 100 \quad (3)$$

Where, PAR_a = PAR above the canopy and PAR_b = PAR below the canopy.

The determination of LAI was done using the inversion of transmitted PAR, as indicated in the equation according to Goudriaan (1988)

$$L = \frac{\left[\left(1 - \frac{1}{2K} \right) f_b - 1 \right] \ln \tau}{A(1 - 0.47 f_b)} \quad (4)$$

Where, L is leaf area index; K is the extinction coefficient for the canopy, given as $k = \frac{1}{2 \cos \theta}$ with θ the zenith angle of the sun; f_b is the fraction of incident PAR; τ is the ration of PAR measured below the canopy to PAR above the canopy; A is given as $A = 0.283 + 0.785a - 0.159a^2$, with a the leaf absorptivity in the PAR band (typically around 0.9).

Table 1: Treatments in the two sites (ATC-Embu and Kamujine)

| Treatment | Cropping system | Treatment | Cropping system |
|-----------|---------------------|-----------|---------------------|
| T1 | Sole maize | T4 | Maize-Soybean (2:2) |
| T2 | Sole soybean | T5 | Maize-Soybean (2:4) |
| T3 | Maize-Soybean (1:1) | T6 | Maize-Soybean (2:6) |

Table 2: Effect of intercropping pattern on maize yields and harvest index during 2012 LR and 2012 SR at Embu and Kamujine sites

| Location | Treatment | Stover yield (t/ha) | | Grain yield (t/ha) | |
|-----------------------|-----------------------|---------------------|------------|--------------------|-----------|
| | | 2012 LR | 2013 SR | 2012 LR | 2013 SR |
| Embu | Sole maize | 11.73 | 8.14 | 4.95 | 4.58 |
| | Maize-Soybean (1M:1S) | 12.64 | 7.74 | 5.49 | 5.16 |
| | Maize-Soybean (2M:2S) | 13.12 | 7.62 | 6.11 | 5.62 |
| | Maize-Soybean (2M:4S) | 8.89 | 4.91 | 4.05 | 3.48 |
| | Maize-Soybean (2M:6S) | 5.26 | 4.45 | 2.74 | 3.16 |
| <i>p-value</i> | | 0.0001*** | <0.0001*** | <0.0001*** | 0.0467* |
| LSD _(0.05) | | 2.59 | 1.31 | 0.87 | 1.79 |
| Kamujine | Sole maize | 3.81 | 6.00 | 3.36 | 2.98 |
| | Maize-Soybean (1M:1S) | 3.87 | 6.34 | 3.09 | 3.44 |
| | Maize-Soybean (2M:2S) | 2.95 | 6.55 | 3.07 | 3.55 |
| | Maize-Soybean (2M:4S) | 2.59 | 4.83 | 2.47 | 2.86 |
| | Maize-Soybean (2M:6S) | 1.90 | 3.97 | 1.9 | 1.82 |
| <i>p-value</i> | | 0.0461* | 0.0005*** | 0.0704ns | 0.0006*** |
| LSD _(0.05) | | 1.43 | 1.00 | 1.07 | 0.64 |

ns – not significant; *significant at $p \leq 0.05$; **significant at $p < 0.01$; ***significant at $p < 0.001$.

Table 3: Effect of intercropping pattern on soybean yields and harvest index during 2012 LR and 2012 SR at Embu and Kamujine sites

| Location | Treatment | Stover yield (t ha ⁻¹) | | Grain yield (t ha ⁻¹) | |
|-----------------------|-----------------------|------------------------------------|-----------|-----------------------------------|-------------|
| | | 2012 LR | 2012 SR | 2012 LR | 2012 SR |
| Embu | Sole soybean | 1.41 | 1.45 | 1.44 | 1.65 |
| | Maize-Soybean (1M:1S) | 0.40 | 0.76 | 0.26 | 0.41 |
| | Maize-Soybean (2M:2S) | 0.53 | 0.55 | 0.35 | 0.42 |
| | Maize-Soybean (2M:4S) | 0.60 | 0.83 | 0.46 | 0.76 |
| | Maize-Soybean (2M:6S) | 1.03 | 1.17 | 0.88 | 1.01 |
| <i>p-value</i> | | 0.0015** | 0.0001*** | 0.0002*** | < 0.0001*** |
| LSD _(0.05) | | 0.43 | 0.28 | 0.39 | 0.18 |
| Kamujine | Sole soybean | 1.09 | 1.27 | 0.56 | 0.72 |
| | Maize-Soybean (1M:1S) | 0.18 | 0.21 | 0.05 | 0.07 |
| | Maize-Soybean (2M:2S) | 0.19 | 0.24 | 0.04 | 0.08 |
| | Maize-Soybean (2M:4S) | 0.52 | 0.58 | 0.17 | 0.30 |
| | Maize-Soybean (2M:6S) | 0.78 | 0.83 | 0.24 | 0.37 |
| <i>p-value</i> | | < 0.0001*** | 0.0004*** | 0.0007*** | <0.0001*** |
| LSD _(0.05) | | 0.13 | 0.39 | 0.20 | 0.08 |

ns – not significant; *significant at $p \leq 0.05$; **significant at $p < 0.01$; ***significant at $p < 0.001$.

DATA ANALYSIS

Data of maize and soybean yields, PAR and LAI were subjected to analysis of variance using SAS version 8. To

test for significant differences between different cropping pattern and conventional intercropping systems, the yields were subjected to *t-student* test at 95 percent of significance level ($p < 0.05$).

Table 4: Shading effect on soybean yield in the 2M:4S during 2012 LR and 2012 SR at Embu and Kamujine sites


| Location | Row position | Maize-Soybean (2M:4S) | | | |
|-----------------------|--------------|-----------------------|----------|--------------------|----------|
| | | Stover yield (t/ha) | | Grain yield (t/ha) | |
| | | 2012 LR | 2012 SR | 2012 LR | 2012 SR |
| Embu | Middle | 0.07 | 0.095a | 0.05 | 0.061a |
| | Closer | 0.06 | 0.086b | 0.04 | 0.052b |
| <i>p-value</i> | | 0.0278* | 0.0113** | 0.0009*** | 0.0108** |
| LSD _(0.05) | | 0.01 | 0.007 | 0.01 | 0.007 |
| Kamujine | Middle | 0.05 | 0.066 | 0.019 | 0.034 |
| | Closer | 0.05 | 0.054 | 0.015 | 0.027 |
| <i>p-value</i> | | 0.7292 | 0.0088** | 0.0251* | 0.0085** |
| LSD _(0.05) | | 0.01 | 0.009 | 0.004 | 0.005 |

ns – not significant; *significant at $p \leq 0.05$; **significant at $p < 0.01$; ***significant at $p < 0.001$.

Table 5: Shading effect on soybean yield in the 2M:6S treatment during 2012 LR and 2012 SR at Embu and Kamujine sites

| Location | Row position | Maize-Soybean (2M:6S) | | | |
|-----------------------|--------------|-----------------------|----------|--------------------|----------|
| | | Stover yield (t/ha) | | Grain yield (t/ha) | |
| | | 2012 LR | 2012 SR | 2012 LR | 2012 SR |
| Embu | Middle | 0.11 | 0.13a | 0.10 | 0.12a |
| | Closer | 0.08 | 0.12a | 0.07 | 0.09b |
| <i>p-value</i> | | 0.0179* | 0.1008 | 0.0236* | 0.0297* |
| LSD _(0.05) | | 0.03 | 0.02 | 0.03 | 0.02 |
| Kamujine | Middle | 0.086 | 0.10a | 0.024 | 0.065a |
| | Closer | 0.066 | 0.060b | 0.016 | 0.036b |
| <i>p-value</i> | | 0.1049 | 0.0051** | 0.0305* | 0.0017** |
| LSD _(0.05) | | 0.025 | 0.03 | 0.007 | 0.02 |

ns – not significant; *significant at $p \leq 0.05$; **significant at $p < 0.01$; ***significant at $p < 0.001$.

**Figure 1:** Relationship between soybean grain yield with PAR intercepted and LAI in middle rows during 2012 SR at Embu site

RESULTS AND DISCUSSION

Effects of maize-soybean intercropping patterns on maize and soybean yields

Maize yields

At Embu during both seasons, maize stover and grain yields were significantly affected by the intercropping

pattern (Table 2). For instance, during 2012 LR the MBILI treatment recorded significantly higher stover and grain yields (13.12 t ha^{-1} , $p=0.0001$ and 6.11 t ha^{-1} , $p < 0.0001$, respectively) than all the other treatments. During the 2012 SR, similar trend emerged with the MBILI treatment recording significantly the highest stover and grain yield 7.62 t ha^{-1} , $p < 0.0001$ and 5.62 t ha^{-1} , $p=0.0467$, respectively (Table 2).

During both seasons at Kamujine site, maize stover




Figure 2: Relationship between soybean stover yield with PAR intercepted and LAI in middle rows during 2012 SR at Kamujine site




Figure 3: Relationship between soybean grain yield with PAR intercepted and LAI in middle rows during 2012 SR at Kamujine site

yield was significantly ($p \leq 0.05$) affected by the intercropping pattern. For instance, during 2012 LR the conventional treatment produced significantly the highest stover yield (3.87 t ha^{-1} , $p=0.0461$). During this season, the grain yield was not significantly affected by the intercropping patterns ($p=0.0704$); however, the sole maize treatment recorded numerically the highest value of 3.36 t ha^{-1} while and 2M:6S treatment recorded the lowest yield. The lower maize yields under 2M:6S treatment could be related to the lower plant density compared to other treatments. During the 2012 SR, the MBILI treatment recorded significantly the highest stover and grain yield (6.55 t ha^{-1} , $p=0.0005$ and 3.55 t ha^{-1} , $p=0.0006$, respectively) than all the other treatments, except sole maize and conventional treatments (Table 2).

Similar results of increased maize yield under MBILI treatment found at Embu site were also reported by Undie *et al.* (2012) in Nigeria under maize-soybean

intercrop, Mucheru-Muna *et al.* (2010) in the central highlands of Kenya and Woomer *et al.* (2004) in western Kenya with maize-beans, and Solanki *et al.* (2011) in India with maize-blackgram. This could be due to the fact that the MBILI intercrop arrangement offer better opportunities to the components to utilize available resources more effectively than the conventional (1M:1S) intercropping pattern. Rajat De and Singh (1979) stated that the modified 2M:2S system affords a better solar energy harvest than the 1M:1S crop arrangement, because the former might have provided sufficient light to the lower leaves to continue photosynthesis. Furthermore, Brintha and Seran (2009) stated that productivity rates increases with LAI because of increased total light interception, but larger LAI values often cause no more increases and then decreases on a ground basis, probably due to respiratory CO_2 loss from heavily shaded leaves and stems. However, when the

Table 6: Effects of intercropping patterns on PAR and LAI of maize and soybean during 2013 SR at Embu and Kamujine sites

| Location | Treatment | Crop | 35 DAP | | 49 DAP | | 63 DAP | | |
|----------|-----------------------|---------|--------|------------|------------|---------|---------|---------|--------|
| | | | PAR% | LAI | PAR% | LAI | PAR% | LAI | |
| Embu | Sole maize | Maize | 57.23 | 1.03 | 66.50 | 1.28 | 83.40 | 3.61 | |
| | Sole soybean | Soybean | 58.23 | 1.09 | 54.97 | 1.00 | 73.12 | 2.87 | |
| | Maize-Soybean (1M:1S) | Maize | 32.41 | 0.47 | 56.55 | 0.99 | 74.19 | 2.74 | |
| | | Soybean | 41.65 | 0.66 | 56.67 | 0.99 | 78.05 | 3.06 | |
| | Maize-Soybean (2M:2S) | Maize | 45.88 | 0.74 | 61.00 | 1.24 | 74.10 | 3.21 | |
| | | Soybean | 53.70 | 0.95 | 66.92 | 1.45 | 84.15 | 4.26 | |
| | Maize-Soybean (2M:4S) | Maize | 42.67 | 0.70 | 55.58 | 0.95 | 69.05 | 2.34 | |
| | | Soybean | 55.89 | 1.00 | 48.74 | 0.87 | 68.24 | 2.40 | |
| | Maize-Soybean (2M:6S) | Maize | 46.34 | 0.75 | 51.38 | 0.87 | 66.95 | 2.23 | |
| | | Soybean | 53.82 | 0.95 | 59.13 | 1.10 | 77.73 | 3.35 | |
| | <i>p</i> – value | | | 0.0002*** | 0.0003*** | 0.2850 | 0.3564 | 0.0310* | 0.0962 |
| | LSD _(0.05) | | | 10.10 | 0.25 | 14.97 | 0.52 | 10.95 | 1.33 |
| Kamujine | Sole maize | Maize | 49.86 | 0.85 | 69.39 | 1.52 | 76.47 | 2.28 | |
| | Sole soybean | Soybean | 62.08 | 1.18 | 75.72 | 1.67 | 80.93 | 2.65 | |
| | Maize-Soybean (1M:1S) | Maize | 36.24 | 0.55 | 67.17 | 1.37 | 78.90 | 2.54 | |
| | | Soybean | 32.51 | 0.48 | 54.79 | 0.95 | 67.68 | 1.89 | |
| | Maize-Soybean (2M:2S) | Maize | 43.71 | 0.71 | 80.69 | 2.06 | 84.23 | 3.24 | |
| | | Soybean | 41.66 | 0.66 | 70.60 | 1.46 | 79.48 | 2.60 | |
| | Maize-Soybean (2M:4S) | Maize | 37.44 | 0.57 | 64.10 | 1.21 | 71.74 | 2.08 | |
| | | Soybean | 41.38 | 0.66 | 65.62 | 1.38 | 73.57 | 2.00 | |
| | Maize-Soybean (2M:6S) | Maize | 48.87 | 0.82 | 67.97 | 1.37 | 76.84 | 2.34 | |
| | | Soybean | 23.26 | 0.33 | 65.17 | 1.37 | 80.98 | 2.48 | |
| | <i>p</i> – value | | | <0.0001*** | <0.0001*** | 0.0476* | 0.0342* | 0.1417 | 0.2026 |
| | LSD _(0.05) | | | 8.83 | 0.20 | 13.35 | 0.54 | 11.05 | 0.93 |

ns – not significant; *significant at $p \leq 0.05$; **significant at $p < 0.01$; ***significant at $p < 0.001$.

plant population is different the yield difference is determined by population density rather than crop arrangement (Rajat De and Singh, 1979), which was not the case in this particular situation. Similarly, Sikirou and Wydra (2008), Dapaah *et al.* (2008), Mpairwe *et al.* (2002) and Olufemi *et al.* (2001) reported higher cereal grain yield under intercropping system compared to its sole crop. Maize grain yield differs with different legume species and intercropping produces higher maize grain yield than in pure stand (Sikirou and Wydra, 2008).

On the other hand, Thobatsi (2009) in South Africa, Silwana and Lucas (2002) also observed reduction in maize yield under intercropping system compared to its monocrop, under similar environment of limited rainfall conditions, as it was observed at Kamujine site during 2012 LR. In this site it was necessary to supplement with irrigation water because the crop had started to suffer from water stress due to lack of rainfall during flowering period, as it can be seen in the Figure 3.3. Amede (1995) stated that one of the factors that reduces maize grain yield is dry conditions that occur specially during the flowering period. Higher populations under intercrops compared to monocrop under stress conditions might result in intercrop yields being lower than sole crop yields due to increased competition for moisture (Natarajan and Willey, 1986). Yield reductions involving one or all intercropping components in intercropping could be

associated to inter-specific competition for nutrients, moisture and/or space (Adaniyan *et al.*, 2007). Moreover, Kamujine has a light texture soil, which has low moisture holding capacity (Jaetzold *et al.*, 2006), resulting therefore in reduced yields under intercropping (Natarajan and Willey, 1986).

The efficiency of a crop variety to convert the dry matter into economic yield is determined by its harvest index. The higher the value, the higher will be dry matter conversion efficiency. The absence of significant differences in HI of maize observed during 2012 SR at Embu and Kamujine sites agrees with results by Haseebur-Rehman *et al.* (2010) and Egbe, Alibo and Nwueze (2010) in maize-cowpea intercropping; Saleem *et al.* (2011) in maize-legume intercropping systems; and, Carruthers *et al.* (2000) in maize-soybean intercropping who reported that the intercropping systems did not affect the harvest index of maize component.

Soybean yields and components

During both seasons in both sites, the soybean yield was significantly affected by the intercropping pattern. During the 2012 LR, the yields were reduced by 60 and 81 percent due to the intercropping with maize, at Embu and Kamujine, respectively; whereas, during the 2012 SR, the




Figure 4: Relation between sobean grain yield under MBILI treatment with PAR and LAI during 2012 SR at Embu site




Figure 5: PAR Intercepted (%) during 2013 SR at Embu (a) and Kamujine (b) sites

yields were reduced by 52 and 78 percent as a result of intercropping with maize, at Embu and Kamujine, respectively (Table 3). At Embu, during 2012 LR, the sole soybean observed significantly the highest stover yield (1.41 t ha^{-1} , $p=0.0015$) than all the intercropping patterns, followed by the 2M:6S intercrop pattern with 1.03 t ha^{-1} which was statistically different from MBILI (0.53 t ha^{-1} , $p = 0.0015$) and conventional (0.40 t ha^{-1} , $p = 0.0015$) intercropping patterns, but not significantly different from the 2M:4S intercropping pattern. Also, the sole soybean treatment gave statistically the highest grain yield (1.44 t ha^{-1} , $p=0.0002$) than all the intercropping patterns, followed again by the 2M:6S treatment (0.88 t ha^{-1} , $p=0.0002$), which was significantly different from other intercropping patterns (conventional, MBILI and 2M:4S). Similar results were obtained during 2012 SR, where sole soybean treatment recorded statistically the highest stover yield (1.45 t ha^{-1} , $p<0.0001$) than all the other treatments, followed by the 2M:6S treatment (1.17 t ha^{-1} , $p<0.0001$), which was statistically different from the rest of the intercropping patterns. Although the conventional treatment observed 28 per cent higher stover yield than the MBILI treatment, they were not statistically different

among them. The monocropped soybean observed also the highest grain yield (1.65 t ha^{-1} , $p<0.0001$) than all other treatments, followed by the 2M:6S treatment with 1.01 t ha^{-1} ($p<0.0001$), which significantly differed from the rest of the treatments. At Kamujine during 2012 LR, the sole soybean treatment gave statistically the highest stover and grain yields (1.09 t ha^{-1} , $p<0.0001$ and 0.56 t ha^{-1} , $p=0.0007$, respectively) than all the other treatments, followed by the 2M:6S treatment with 0.78 t ha^{-1} for the stover yield, which was significantly different ($p<0.0001$) from the other intercropping patterns. During the 2012 LR season, there was significant positive correlation between the soybean grain yield and, and soil mineral N, with $r=0.73$ ($p=0.0002$) and $r=0.76$ ($p<0.0001$), respectively. During 2012 SR, monocropped soybean treatment observed also significantly the highest stover yield (1.27 t ha^{-1} , $p=0.0004$) than all the other treatments. Similarly, sole soybean treatment recorded the highest grain yield of 0.72 t ha^{-1} ($p<0.0001$) than all the other treatments. In general, the yields increased with about 23 and 26 percent in the second season compared to the first season, at Embu and Kamujine, respectively. Probably due to the cumulative effects of the goat

Table 7: Effects of maize shade on PAR intercepted of soybean during 2013 SR at Embu and Kamujine sites

| Location | Row position | 35 DAP | | 49 DAP | | 63 DAP | |
|-----------------------|--------------|------------|------------|------------|----------|-----------|------------|
| | | 2M:4S | 2M:6S | 2M:4S | 2M:6S | 2M:4S | 2M:6S |
| | | PAR | PAR | PAR | PAR | PAR | PAR |
| Embu | Middle | 63.94 | 61.71 | 66.87 | 68.68 | 78.03 | 87.05 |
| | Closer | 43.97 | 42.71 | 47.75 | 49.23 | 66.05 | 71.25 |
| <i>p</i> – value | | 0.001*** | <0.0001*** | <0.0001*** | 0.0029** | 0.0001*** | 0.0092** |
| LSD _(0.05) | | 9.13 | 6.16 | 7.12 | 11.58 | 5.59 | 11.24 |
| Kamujine | Middle | 48.82 | 53.23 | 61.69 | 69.08 | 75.55 | 84.51 |
| | Closer | 31.61 | 46.44 | 47.11 | 49.18 | 63.15 | 63.41 |
| <i>p</i> – value | | <0.0001*** | 0.012* | 0.0051** | 0.0021** | 0.0027** | <0.0001*** |
| LSD _(0.05) | | 5.31 | 5.05 | 9.85 | 11.36 | 7.74 | 6.53 |

ns – not significant; *significant at $p \leq 0.05$; **significant at $p < 0.01$; ***significant at $p < 0.001$.

Table 8: Effects of maize shade on LAI of soybean during 2013 SR at Embu and Kamujine sites

| Location | Row position | 35 DAP | | 49 DAP | | 63 DAP | |
|-----------------------|--------------|------------|---------|--------|--------|--------|---------|
| | | 2M:4S | 2M:6S | 2M:4S | 2M:6S | 2M:4S | 2M:6S |
| | | LAI | LAI | LAI | LAI | LAI | LAI |
| Embu | Middle | 1.24 | 1.17 | 1.33 | 1.38 | 3.23 | 4.20 |
| | Closer | 0.71 | 0.67 | 0.81 | 0.83 | 2.19 | 2.73 |
| <i>p</i> – value | | 0.001** | 0.001** | <0.001 | 0.0028 | 0.001 | 0.0078 |
| LSD _(0.05) | | 0.25 | 0.20 | 0.17 | 0.33 | 0.58 | 1.02 |
| Kamujine | Middle | 0.82 | 0.93 | 1.17 | 1.40 | 2.43 | 3.04 |
| | Closer | 0.46 | 0.76 | 0.80 | 0.83 | 1.62 | 1.62 |
| <i>p</i> – value | | <0.0001*** | 0.0124 | 0.008 | 0.0017 | 0.001 | <0.0001 |
| LSD _(0.05) | | 0.11 | 0.13 | 0.27 | 0.32 | 0.45 | 0.43 |

ns – not significant; *significant at $p \leq 0.05$; **significant at $p < 0.01$; ***significant at $p < 0.001$.

manure that was applied in the first and second seasons, and also due to root and nodule senescence.

At Embu during the 2012 LR, there were significant differences in soybean stover yield ($p=0.0278$) among the rows in 2M:4S treatment, where the row next to the maize plant observed 16.7 percent lower yield than the row far away from the maize plant. For the grain yield, the row far away from the maize plant gave 20 percent significantly higher ($p=0.0009$) yield than the row next to maize plant. During the 2012 SR, the soybean stover and grain yields were 9.5 and 14.8 percent significantly ($p=0.0113$ and $p=0.0108$, respectively) higher in the middle row than in the row next to the maize plant, respectively. At Kamujine during the 2012 LR, there were no significant differences ($p=0.7292$) on soybean stover yield between the middle row and the row next to the maize plant; whereas the grain yield in the middle row was 21 percent significantly ($p=0.0251$) higher than in the row next to the maize plant. During the 2012 SR, the soybean stover and grain yields were 22.2 and 25.9 percent significantly ($p=0.0088$ and $p=0.0085$,

respectively) lower in the closer row than in the middle row, respectively (Table 4).

At Embu during the 2012 LR, there were significant differences in soybean stover yield ($p=0.0179$) among rows in 2M:6S intercropping pattern. The middle row recorded about 27.3 percent higher yield than the row next to the maize row. For the grain yield, the middle row observed 30 per cent significantly higher ($p=0.0236$) than the row closer to maize plant. During the 2012 SR, the soybean stover yield did not show significant differences ($p=0.1008$) between the row next to the maize plant and one in the middle row; however, it was numerically 7.7 percent higher in the middle row than in the row next to the maize plant. The grain yield was 25 percent significantly ($p=0.0297$) higher in middle row than in the closer row. At Kamujine during the 2012 LR, there were no significant differences ($p=0.1049$) on soybean stover yield between the row middle row and the row closer to the maize plant; whereas the grain yield in the middle row was 33.3 per cent significantly ($p=0.0305$) higher than in the closer row. During the 2012 SR, the soybean stover

and grain yields were 40.0 and 44.6 percent significantly ($p=0.0051$ and $p=0.0017$, respectively) higher in the middle row than in the closer row, respectively (Table 5).

At Embu site, during the 2012 SR there was statistically significant strong positive correlation between soybean grain yields of the middle row and PAR intercepted ($r=0.998$; $p=0.0019$) and LAI ($r=0.992$; $p=0.0078$), at 49 DAP (Figure 1).

At Kamujine site, during 2012 SR also middle rows showed existence of significant correlation between soybean yields and light intercepted and Leaf area index, at 63 DAP. For instance, soybean stover yield was strongly positive correlated with the PAR intercepted ($r=0.95$; $p=0.0463$) and LAI ($r=0.99$; $p=0.0088$), as indicated by the Figure 2. Similarly, there was strong positive correlation between soybean grain yield with the amount of light intercepted ($r=0.997$; $p=0.0027$) and LAI ($r=0.99$; $p=0.0085$) (Figure 3)

Effects of maize-soybean intercropping patterns on light interception and Leaf Area Index

At Embu during the 2012 SR, significant differences ($p<0.001$ and $p<0.05$) were observed in light interception (PAR) and leaf area index (LAI) for maize and soybean during the sampling period as affected by the intercropping patterns, except during 49 days after planting (DAP). For instance, during the 35 DAP sole soybean intercepted significantly ($p=0.0002$) more light (58.23 percent) and leaf area index (LAI) of 1.03 ($p=0.0003$) than all other treatments and/or crop, excluding sole maize, soybean in 2M:2S, 2M:4S and in 2M:6S treatments. In this period only soybean under MBILI treatment showed existence of significant strong correlation between grain yield and PAR intercepted ($r=0.98$; $p=0.0254$) and LAI ($r=0.97$; $p=0.0324$), as shown by the figure 4.6. While at 63 DAP the soybean in MBILI treatment observed statistically ($p=0.0310$) the highest light interception (84.15 percent) than sole soybean, maize and soybean under 2M:4S, and maize under 2M:6S treatments (Table 6). At Kamujine site during the same sampling period, where both PAR and LAI were significantly (35 DAP with $p<0.0001$ and $p<0.0001$; and, 49 DAP with $p=0.0476$ and $p=0.0342$, respectively) affected by the intercropping patterns. At 35 DAP the sole soybean recorded statistically the highest PAR of 62.08 per cent ($p<0.0001$) and LAI of 1.18 ($p<0.0001$) compared to all other treatments. Whereas, during the time of 49 DAP maize under MBILI treatment intercepted significantly more light (80.69 per cent, $p=0.0476$) than all other treatments, excluding sole soybean, sole maize, soybean under MBILI and maize under 2M:6S treatments. In the same period (49 DAP), still maize under MBILI treatment showed the highest LAI of 2.06 than all other treatments, except soybean under monocrop (Table 6).

The higher light interception showed by sole soybean treatment at Embu and Kamujine sites was also reported by Ghanbari, Dahmardeh, Siahsar and Ramroudi (2010), who observed that the cowpea-maize intercropping and maize sole crop had showed a lower light interception compared to cowpea sole crop. Studying wheat-faba bean intercropping, Eskandari (2011) found that the mean of PAR interception averaged over sampling dates by sole cropped bean was significantly higher than that for sole cropped wheat. Studying maize-peas intercropping, Kanton and Dennett (2008) found that at 36 DAP the sole peas were intercepting 30 per cent more light than sole maize. Studying maize-bean intercropping, Tsubo *et al.* (2001) found that sole beans showed higher PAR intercepted than sole maize. Studying southern pea-sweet corn intercropping system, Francis and Decoteau (1993) found that monocropped southern pea had intercepted more light than monocropped sweet corn 52 and 66 DAP. Solanki *et al.* (2011) reported that sole soybean intercepted more light as compared to maize+soybean and sole maize at 75 DAP.

Varies studies, particularly in tropical, have shown that intercrops intercept more PAR than sole crops, for example, Sivakumar and Virmani (1980) with maize-pigeon pea, Bandy-opadhyay (1988) with sorghum-mung bean, and Tsubo *et al.* (2001) with maize-bean intercrops. The present study confirmed that MBILI treatment capture more radiant energy than sole crops, but the differences between this treatment and the sole maize crop (at Embu site) or soybean crop (at Kamujine site) were relatively small and not significant. This might be probably due to the differences in agro-ecological zonation and rainfall patterns between the two sites, where at Embu site (upper midland) had received more rainfall which made the maize crop to respond positively as it allowed faster leaf area growth and plant height, and therefore more light was intercepted; whereas, at Kamujine site (lower midland) had received less rainfall which made the maize crop to retard its initial growth rate compared to soybean and not being able to cover the soil surface very early resulting in lower interception of PAR compared to soybean. Also, Soybean plants, being deep rooted, can to some extent withstand dry periods (Prasad and Brook, 2005).

Environmental factors such as limited moisture, influences the leaf area development (Afuakwa and Crookston, 1984). Tsubo *et al.* (2001) stated that faster leaf area growth can cause higher PAR intercepted during the vegetative stage. Radiation interception varies from seedling emergence to crop harvest (Natarajan and Willey, 1980; Sivakumar and Virmani, 1980) and depends largely on the canopy leaf area (Tsubo *et al.*, 2001). The higher PAR conversion efficiencies of intercrop systems relative to the sole crops may be due to spread of light over greater leaf area, and more efficient distribution of light in their canopies during early stages of growth (Addo-Quaye *et al.*, 2011). Keating and Carberry (1993)

concluded that cereal and legume can differ in PAR interception because of differences in their vertical arrangement of foliage and canopy architecture and can therefore intercept more PAR compared to sole crops. Moreover, it has been found by several investigators (Reddy and Willey, 1981; Sivakumar and Virmani, 1984; Watiki *et al.*, 1993) that LAI patterns follow the patterns of radiation interception.

The higher LAI observed during 49 DAP with intercropped maize at Kamujine agrees with the results of Baldé *et al.* (2011) who found that maize intercropped with pigeonpea or brachiria had higher LAI than sole maize crop; and, Thobatsi (2009) who found maize intercropped with cowpea long duration cultivar had significantly higher LAI of 2.23 compared to medium season cultivars. Higher maize LAI when intercropped with soybean during 2013 SR could have been attributed to sufficient rainfall at the beginning of the season that stimulated maize leaf growth.

Similarly, the reductions of maize LAI under intercropping systems in both sites (Embu and Kamujine) have been also reported by other researchers (Thobatsi, 2009; Zegada-Lizarazu, Izumi and Ijima, 2006; Bilalis *et al.*, 2005). The absence of differences in maize LAI between all treatments in both sites (Embu and Kamujine) at 63 DAP and at 49 DAP in Embu site correspond with results of Filho (2000) who did not find any significant differences between sole maize and maize intercropped with cowpea. Twala and Ossom (2004) also did not find any significant differences in LAI between maize monocrop and maize intercropped with sugar beans or groundnuts. These results support the fact that different climatic conditions will result in different leaf development patterns of maize, with or without intercropping.

In general, at Embu site monocropped maize and the MBILI treatments intercepted numerically more light than all other treatments, with averages of 69.04 and 64.29 percent during the sampling period, respectively. The MBILI treatment intercepted 10.43 per cent more light than the conventional treatment (Figure 5). While, at Kamujine site monocropped soybean and the MBILI treatment intercepted numerically more light than all other treatments, with averages of 72.91 and 66.73 percent, respectively. The PAR intercepted by the MBILI treatment was 15.66 percent above the one that the conventional treatment had intercepted (Figure 5).

Similar results were also reported by Woomer and Tungani (2003) who found that beans had intercepted 20 per cent more light under MBILI treatment than under conventional crop arrangement. These findings demonstrate the advantages of light penetration to the MBILI intercropping pattern which offer realizable potential for smallholder farmers to improve the performance of their maize-legume enterprise (Woomer and Tungani, 2003).

The Table 7 shows that in both sites (Embu and

Kamujine) the amount of light intercepted by the legume component and LAI were significantly ($p < 0.05$) affected by maize shade. For instance at Embu site, 35 DAP the middle row intercepted significantly ($p = 0.001$ and $p < 0.0001$) 19.97 and 19.0 per cent more light than the closer row, in the 2M:4S and 2M:6S treatments, respectively. During the time of 49 DAP, the closer row had significantly ($p < 0.0001$ and $p = 0.0029$) intercepted 19.12 and 19.45 percent less light than the middle row, in the 2M:4S and 2M:6S treatments, respectively. And at 63 DAP, the PAR intercepted by the middle row was 11.98 and 15.80 percent statistically ($p = 0.0001$ and $p = 0.0029$) higher than the one that was intercepted by the closer row, in the 2M:4S and 2M:6S treatments, respectively. Similarly, at Kamujine the closer row had significantly ($p < 0.0001$ and $p = 0.012$) intercepted 17.21 and 6.79 percent less light than the middle row, in the 2M:4S and 2M:6S treatments, respectively, at the time of 35 DAP. During the time of 49 DAP, the middle row had significantly ($p = 0.0051$ and $p = 0.0021$) intercepted 14.58 and 19.90 percent more light than the closer row, in the 2M:4S and 2M:6S treatments, respectively. At 63 DAP, the PAR intercepted by the closer row was 12.40 and 21.10 percent statistically ($p = 0.0027$ and $p < 0.0001$) lower than the one that was intercepted by the middle row, in the 2M:4S and 2M:6S treatments, respectively (Table 7).

At Embu site during 2013 SR, at 35 DAP the middle row observed significantly ($p = 0.001$ and $p < 0.001$) 1.24 and 1.17 higher LAI than the closer row with 0.71 and 0.67, in the 2M:4S and 2M:6S treatments, respectively. During the time of 49 DAP, the closer row had significantly ($p < 0.0001$ and $p = 0.0029$) recorded 0.52 and 0.55 lower LAI than the middle row, in the 2M:4S and 2M:6S treatments, respectively. At 63 DAP, the LAI observed by the middle row was 1.04 and 1.47 statistically ($p = 0.001$ and $p = 0.0078$) higher than the one that was recorded by the closer row, in the 2M:4S and 2M:6S treatments, respectively (Table 4.17).

At Kamujine site, during 35 DAP the closer row had significantly ($p < 0.0001$ and $p = 0.0124$) observed lower LAI with values of 0.46 and 0.76 than the middle row (0.82 and 0.93), in the 2M:4S and 2M:6S treatments, respectively. During the time of 49 DAP, the middle row had significantly ($p = 0.008$ and $p = 0.0017$) recorded higher LAI with values of 1.17 and 1.40 than the closer row that observed 0.80 and 0.83, in the 2M:4S and 2M:6S treatments, respectively. At 63 DAP, the LAI recorded by the closer row was 0.81 and 1.42 statistically ($p = 0.001$ and $p < 0.0001$) lower than the one that was showed by the middle row, in the 2M:4S and 2M:6S treatments, respectively (Table 4.13). Similar effects of shading on legume component have been reported by several authors (Hang *et al.*, 1984; Stirling *et al.*, 1990; Ali, Jeffers and Henderlong (2003). For instance, Ali *et al.* (2003) found that alternate row of soybean had intercepted 31 percent more PAR than alternate narrow rows at 74 DAP.

The reduction of soybean yields under intercropping with maize or sorghum has also been reported by several researchers (Ijoyah and Fanen, 2012; Muoneke *et al.*, 2007; Muneer *et al.*, 2004; Heibsch *et al.*, 1995; Olufajo, 1992; Pal, Oseni, and Norman, 1992; Neupane, 1983). This reduction in soybean yields under intercropping could be due to interspecific competition between the intercrop components for water, light, air and nutrients, and also the aggressive effects maize (C_4 species) on soybean, a C_3 species (Muoneke *et al.*, 2007). According to Heibsch *et al.* (1995), crops with C_4 photosynthetic pathways have been known to be dominant when intercropped with C_3 species like soybean. The shading of soybean by the maize plants (taller) may also have contributed to the reduction of the yields of intercropped soybean. Olufajo (1992) reported that shading by the taller plants in mixture could reduce the photosynthetic rate of the lower growing plants and thereby reduce their yields. This observation was confirmed in this study where lower soybean yields were recorded in the rows next to the maize plant compared to the middle row, and the yields positively correlated with the PAR intercepted. In addition, Lesoing and Francis (1999) stated that water stress and shading were probably the two major factors, which contribute to reduced legume component yield under intercropping. Moreover, Natarajan and Willey (1980) and Sivakumar and Virmani (1980) reported that stover yield often shows a positive correlation with the amount of radiation intercepted by crops in intercropping systems. However in the current this study there was a positive correlation between grain yield with the amount of radiation intercepted by crops in intercropping treatments.

The reduction of harvest index of the legume component observed in this study was also reported by other researchers (Alhassan, Kalu and Egbe, 2012). The reduction was mainly due to maize shading effects on soybean, which caused the legume component to allocate its photosynthates to vegetative growth and height increasing for competing with taller maize. On the other hand, the findings of this study at Kamujine site are in agreement with Carruthers *et al.* (2000) who reported that the HI for intercrop soybean was not significantly different from monocrop soybean.

CONCLUSIONS

The maize-soybean intercropping patterns had significant effect on maize stover and grain yields during both seasons and sites. The MBILI treatment recorded significantly the highest maize stover and grain yields. During both seasons in both sites, the soybean yield was significantly affected by the intercropping pattern. During the 2012 LR, the yields were reduced by 60 and 81 percent and during the 2012 SR, the yields were reduced by 52 and 78 percent as a result of intercropping with

maize. The intercropping patterns affected significantly the PAR intercepted and the leaf area index at both sites. The soybean sole crop intercepted significantly more light and leaf area index (LAI) than all other treatments and/or crop. Towards the end of the season, the soybean in MBILI treatment observed statistically the highest light interception than other treatments.

REFERENCES

- Addo-Quaye AA, Darkwa AA, Ocloo GK (2011). Yield and Productivity of Component Crops in a Maize-Soybean Intercropping System as Affected By Time of Planting and Spatial Arrangement. *Journal of Agricultural and Biological Science*. Vol. 6, No. 9: 50-57.
- Adeniyi ON, Akande SR, Balogun MO, Saka JO (2007). Evaluation of Crop Yield of African Yam Bean, Maize and Kenaf under Intercropping Systems. *American – Eurasian J. Agric. & Environ. Sci.* 2(1): 99 – 102.
- Afuakwa JJ, Crookston NR (1984). Using the Kernel Milk Line to Visually Monitor Grain Maturity in Maize. *Crop Sci.* 24(4), 687-691.
- Alhassan GA, Kalu BA, Egbe OM (2012). Influence of Planting Densities on the Performance of Intercropped Bamabra Groundnut with Cowpea in Makurdi, Benue State, Nigeria. *International Journal of Development and Sustainability*. Vol. 1(3):1-20.
- Ali M, Jeffers DL, Henderlong PR (2003). Interrelationship Between Leaf Area, Light Interception and Growth Rate in a Soybean – Wheat System. *Asian Journal of Plant Science* 2(8): 605-612.
- Amede T (1995). Yield Gain and Risk Minimization in Maize Through Cultivar Mixtures in Semi-Arid Zones of the Rift Valley in Ethiopia. *Exp. Agric.* 31: 161 – 168.
- Baldé AB, Scopel E, Affholder F, Corbeels M, Da Silva FAM, Xavier JHV, Wery J (2011). Agronomic Performance of No-Tillage Relay Intercropping with Maize under Smallholder Conditions in Central Brazil. *Field Crops Research*. 124, 240-251.
- Bambrick A (2009). Soil Organic Carbon in Tree-Based Intercropping Systems of Quebec and Ontario, Canada. MSc. Thesis. McGill University, Montréal, Canada. 93pp.
- Bandy-opadhyay SK (1988). Solar Radiation Interception, Soil Water and Nitrogen Use in Sorghum Based Intercropping System with Legumes. *Indian J. Agron.* 33: 31-36.
- Beets WC (1982). Multiple Cropping and Tropical Farming Systems. Westview Press, Boulder, 156pp.
- Bilalis DJ, Sidiras N, Kakampouki I, Efthimiadou A, Papatheohari Y, Thomopoulos P (2005). Effects of Organic Fertilizer on Maize/Legume Intercrop in a Clay Loam Soil and Mediterranean Climate – Can the Land Equivalent Ratio Index (LER) be Used for Root Development? *J. Food, Agric. & Environ.* 3 (3&4): 117-123.
- Brintha I, and Seran TH (2009). Effect of Paired Row Planting of Radish (*Raphanussativus* L.) Intercropped with Vegetable Amaranthus (*Amaranthus tricolor* L.) on Yield Components of Radish in Sandy Regosol. *J. Agric. Sci.*, 4: 19-28.
- Carruthers K, Prithviraj B, Fe Q, Cloutier D, Martin RC, Smith DL (2000). Intercropping Corn with Soybean, Lupin and Forages: Yield Component Responses. *European Journal of Agronomy*. 12:103-115.
- Dapaah HKm, Asafu-agyei JN, Ennin SA, Yamoah C (2008). Yield Stability of Cassava, Maize, Soybean and Cowpea Intercrops. *J. Agric. Sci.* 140: 73 – 82.
- Egbe OM, Alibo SE, Nwueze I (2010). Evaluation of Some Extra-Early- and Early-Maturing Cowpea Varieties for Intercropping with Maize in Southern Guinea Savanna of Nigeria. *Agric. Biol. J. N. Am.* 1(5): 845-858.
- Ennin SA, Clegg MD, Francis CA (2002). Resource Utilization in Soybean/Maize Intercrops. *African Crop Science Journal*, Vol. 10. No. 3, 2002, pp. 251-261.
- Filho JML (2000). Physiological Responses of Maize and Cowpea to Intercropping. *Pesq. Brasilia*. 35(5), 915-921.
- Francis R, and Decoteau DR (1993). Developing Effective Southern pea and Sweet Corn Intercrop System. *Hort Technology*, Vol. 3(2): 178-

- 184.
- Ghanbari A, Dahmardeh M, Siah SAR BA, Ramroudi M (2010). Effect of Maize (*Zea mays* L.) – Cowpea (*Vigna unguiculata* L.) Intercropping on Light Distribution, Soil Temperature and Soil Moisture in Arid Environment. *Journal of Food, Agriculture & Environment* Vol. 8(1):102-108.
- Goudriaan J (1988). Crop Micrometeorology: A Simulation Study, Centre of Agriculture Publication Documentation, Wageningen, The Netherlands.
- Hang AN, McCloud DE, Boote KJ, Duncan WG (1984). Shade Effects on Growth, Partitioning, and Yield Components of Peanuts. *Crop Sci.* 24, 109-115.
- Haseeb-ur R, Ali A, Muhammad W, Tanveer A, Tahir M, Nadeem MA, Zamir MS (2010). Impact of Nitrogen Application on Growth and Yield of Maize (*Zea mays* L.) Grown Alone and in Combination with Cowpea (*Vigna unguiculata* L.). *American-Eurasian J. Agric. & Environ. Sci.* 7(1):43-47.
- Hiebsch C, Tetio-Kegho F, Chiremba FP (1995). Plant Density and Soybean Maturity in a Soybean-Maize Intercrop. *Agron. J.* 87, 965 – 970.
- Horwith B (1985). A Role for Intercropping in Modern Agriculture. *Biological Sciences* 35(5): 286-291.
- Ijoyah MO, Fanen FT (2012). Effects of Different Cropping Patterns on Performance of maize-soybean Mixture in Makurdi, Nigeria. *Scientific Journal of Crop Science* Vol. 1 (2): 39 – 47.
- Jaetzold R, Schmidt H, Hornetz B, Shisanya CA (2006). Farm Management Handbook of Kenya. Natural Conditions and Farm Information, 2nd edition, vol. 11/C. Ministry of Agriculture/GTZ, Nairobi, Kenya (Eastern Province).
- Jarenyama P, Hesterman OB, Waddington SR, Harwood RR (2000). Relay-Intercropping of Sunnhemp and Cowpea into a Smallholder Maize System in Zimbabwe. *Agron. J.*, 92: 239–244.
- Kanton RAL, Dennett MD (2008). Radiation Capture and Use as Affected by Morphologically Contrasting Maize/Pea in Sole and Intercropping. *West African Journal of Applied Ecology – Volume 13.*
- Keating BA, Carberry PS (1993). Resource Capture and Use in Intercropping: Solar Radiation. *Field Crops Res.* 34: 273-301.
- Lesoing GW, Francis CA (1999). Strip Intercropping of Grain Sorghum/Soybean in Irrigated and Rainfed Environments. *Journal of Production Agriculture*. Vol. 12, no. 4: 601 – 606.
- Liu B, Liu XB, Wang C, Jin J, Harbert SJ, Hashemi M (2010). Response of soybean yield and yield components to light enrichment and planting density. *International Journal of Plant Production*, 4 (1): 1-10.
- Mashingaidze N (2004). The effect of Detassiling and Leaf Stripping on productivity and Weed Pressure in a Maize-Bean intercrop in Chinyika Resettlement Area. MSc. Thesis, Crop Science Department, University of Zimbabwe. 88pgs.
- Mpairwe DR, Sabiiti EN, Ummana NN, Tegegne A, Osuju P (2002). Effects of Intercropping Cereal Crops with Forage Legumes and Source of Nutrients on Cereal Grain Yield and Fodder Dry Matter Yields. *Afri. Crop Sci. J.* 10: 81 – 97.
- Mucheru-Muna M, Pypers P, Mugendi D, Kung'u J, Mugwe J, Merckx R, Vanlauwe B (2010). Staggered Maize–Legume Intercrop Arrangement Robustly Increases Crop Yields and Economic Returns in the Highlands of Central Kenya. *Field Crops Research* 115 (2010) 132–139. 2009 Elsevier B.V. All rights reserved. Journal homepage: www.elsevier.com/locate/fcr.
- Muneer AP, Fida HM, Mumtaz AK, Muhammad IS (2004). Performance of Maize in Intercropping System with Soybean under Different Patterns and Nitrogen Levels. *J. App. Sci.* 4(2), 201-205.
- Muoneke CO, Ogwuche MAO, Kalu BA (2007). Effect of Maize Planting Density on the Performance of Maize/Soybean Intercropping System in a Guinea Savannah Agro-ecosystem. *African Journal of Agricultural Research* Vol. 2 (12), pp. 667-677, December 2007.
- Natarajan M, Willey RW (1980). Sorghum-Pigeonpea Intercropping and the Effects of Plant Population Density. 2. Resource Use. *J. Agric. Sci. Camb.* 95, 59-65.
- Natarajan M, Willey RW (1986). The Effects of Water Stress on Yield Advantages of Intercropping Systems. *Field Crops Res.* 13: 117 – 131.
- Neupane RK (1983). Effects of Planting Arrangement on Maize and Soybean Intercropping. *J. Inst. Agric. Anim. Sci.* 4: 1 – 5.
- Odendo M, Bationo A, Kimani S (2011). Socio-Economic Contribution of Legumes to Livelihoods in Sub-Saharan Africa. In: A. Bationo et al. (eds.), *Fighting Poverty in Sub-Saharan Africa: The Multiple Roles of Legumes in Integrated Soil Fertility Management*. DOI 10.1007/978-94-007-1536-3_3, Springer Science+Business Media B.V. 2011, pp 27-46.
- Ofori, F., and Stern, W. R. (1987). Cereal-legume intercropping systems. *Advances in Agronomy* 40:41-90.
- Olufajo OO (1992). Response of soybean to intercropping with maize in a subhumid tropical environment. *Trop. oil seeds J.* 1, 27-33.
- Olufemi O, Pitan R, Odebiyi JA (2001). The Effects of Intercropping with Maize on the Level of Infestation and Damage of Pod – Sucking Bugs in the Cowpeas. *Crop Protection.* 20: 367 – 372.
- Pal UR, Oseni TO, Norman JC (1992). Effect of Component Densities on Productivity of Soybean/Maize Soybean/Sorghum Intercrop. *J. Agron. Crop Sci.*, 170: 66 – 70.
- Prasad RB, and Brook RM (2005). Effect of Varying Maize Densities on Intercropped Maize and Soybean in Nepal. *Experimental Agriculture* 41:365–382.
- Purcell LC (2000). Soybean Canopy Coverage and Light Interception Measurements Using Digital Imagery. *Crop Sci.* 40: 834-837.
- Rajat De, and Singh SP (1979). Management Practices for Intercropping Systems. Pgs: 17 – 21. In: Proceedings of the International Workshop on Intercropping, 10 -13 Jan 1979, Hyderabad, India. Edited by Willey, R. W. and Garver, C.
- Reddy MS, Willey RW (1981). Growth and Resource Use Studies in an Intercrop of Pearl Millet/Groundnut. *Field Crops Res.* 4, 13-24.
- Reddy MS, Floyd CN, Willey RW (1980) Groundnut in Intercropping Systems. In: Proceedings of the International Workshop on Groundnuts, 13-17 October 1980, ICRISAT Center Patancheru, India.
- Saleem R, Ahmed ZI, Ashraf M, Arif M, Malik MA, Munir M, Khan MA (2011). Response of Maize-Legume Intercropping System to Different Fertility Sources under Rainfed Conditions. *Sarhad J. Agric.* Vol. 27 No. 4: 503-511.
- Sanginga N, Woomer PL (2009). Integrated Soil Fertility Management in Africa: Principles, Practices and Development Process. (eds.). Tropical Soil Biology and Fertility Institute of the International Centre for Tropical Agriculture. Nairobi. 263 pp.
- Schou JB, Jeffers DL, Streeter JG (1978). Effects of Reflectors, Black Boards, or Shades Applied at Different Stages of Plant Development on Yield of Soybeans. *Crop Sci.* 18: 29 – 34.
- Seran TH, Brintha I (2010). Review on Maize Based Intercropping. *Journal of Agronomy* 9 (3): 135–145.
- Sikirou S, and Wydra K (2008). Effect of Intercropping Cowpea with Maize or Cassava on Cowpea Bacterial Blight and Yield. *Journal of Plant Diseases and Protection.* 114 (4): 145 – 151.
- Silwana TT, Lucas EO (2002). The Effect of Planting Combinations and Weeding and Yield of Component Crops of Maize/Bean and Maize/Pumpkin Intercrops. *J. Agric. Sci.* 138: 193 – 200.
- Sivakumar MVK, and Virmani SM (1980). Growth and Resource Use of Maize, Pigeonpea and Maize/Pigeonpea Intercrop in an Operational Research Watershed. *Exp. Agric.* 16, 377-386.
- Sivakumar MVK, and Virmani SM (1984). Crop Productivity in Relation to Interception of Photosynthetically Active Radiation. *Agric. For. Meteorol.* 31, 131-141.
- Solanki NS, Singh D, Sumeriya HK (2011). Resources Utilization in Maize (*Zea mays*)-based Intercropping System under Rainfed Condition. *Indian Journal of Agricultural Science.* 81(6):511-515.
- Steiner KG (1982). Intercropping in Tropical Smallholder Agriculture with Special Reference to West Africa. *Schriftenreihe der GTZ, N. 137*, Eischbon, Germany.
- Stirling CM, Williams JH, Black CR, Ong CK (1990). The Effect of Timing of Shade on Development, Dry Matter Production and Light Use Efficiency in Groundnut (*Arachis hypogaea* L.) under Field Conditions. *Aust. J. Agric. Res.* 41, 633-644.
- Thobatsi T (2009). Growth and Yield Responses of Maize (*Zea mays* L.) and Cowpea (*Vigna unguiculata* L.) in an Intercropping System. Msc. Thesis. University of Pretoria. 159p.
- Tsubo M, Walker S, Mukhala E (2001). Comparison of Radiation Use Efficiency of Mono-/Inter-Cropping Systems with Different Row orientations. *Field Crop Research.* 71: 17 – 29.

- Twala MG, and Ossom EM (2004). Legume-Maize Association Influences Crops Characteristics and Yields. 4th International Crop Science Congress. 26-Set – 01-Oct, 2004. Brisbane, Australia.
- Undie UL, Uwah DF, Attoe EE (2012). Growth and Development of Late Season Maize/Soybean Intercropping in Response to Nitrogen and Crop Arrangement in the Forest Agro-ecology of South Southern Nigeria. *Inter. J. Agric. Res.*,7 (1): 1 – 16.
- Watiki JM, Furai S, Banda JA, Keating BA (1993). Radiation Interception and Growth of Maize/Cowpea intercrop as affected by Plant Density and Cowpea Cultivar. *Field Crops Research*. 35, 123-133.
- Willey RW, Natarajan M, Reddy MS, Rao MR, Nambiar PTC M, Kannaiyan J, Bhatnagar VS (1983). Intercropping Studies with Annual Crops. In: Better Crop for Food, Nugent, J. and O'Connor M. (eds). Pitman Co., London.
- Willey RW, Rao MR, Natarajan M (1980). Traditional Cropping Systems with Pigeon pea and Their Improvement. In: Nene, Y. L. and Kumble, V. ICRISAT. 1981. Proceedings of the International Workshop on Pigeon peas, Volume 1, 15-19 December 1980, Patancheru, A.P., India.
- Woomer PL, Tungani JO (2003). Light Availability within an Innovative Maize-Legume Intercropping System in Western Kenya. African Crop Science Conference Proceedings, Vol. 6:42-46.
- Woomer PL, Langat M, Tungani JO (2004). Innovative maize-legume intercropping results in above- and below-ground competitive advantages for understory legumes. *West African Journal of Applied Ecology* 6, 85-94.
- Xiaolei S, Zhifeng W (2002). The optimal Leaf Area Index for Cucumber Photosynthesis and Production in Plastic Green House. ISHS Acta Horticulturae, 633. XXVI International Horticultural Congress.
- Xinyou YE, Schapendonk ADH CM, Xuhua Z (2003). Some Quantitative Relationships Between Leaf Area Index and Canopy Nitrogen Content and Distribution. *Annals of Botany*, 9:893-903.
- Zegada-Lizarazu W, Izumi Y, Iijima M (2006). Water Competition of Intercropped Pearl Millet with Cowpea under Drought and Soil Compaction Stress. *Plant Prod. Sci.* 9(2), 123-132.