



Global Advanced Research Journal of Environmental Science and Toxicology (ISSN: 2315-5140) Vol.2(1) pp. 011-021, January, 2013
Available online <http://garj.org/garjest/index.htm>
Copyright © 2013 Global Advanced Research Journals

Full Length Research Paper

Protective effects of tomato paste and Vitamin E on Atrazine-Induced Hematological and biochemical characteristics of *Clarias gariepinus* (Burchell, 1822)

Imam A. A. Mekkawy^{1,2}, Usama M. Mahmoud² and Rehab H. Mohammed²

¹Biology Department, Faculty of Science, Taif University, Saudi Arabia, iaa_mekkawy@yahoo.co.uk

²Department of Zoology, Faculty of Science, Assiut University, Egypt.

Accepted December 11, 2012

The pesticides have different adverse impacts on different life stages of fish species with attempts to use dietary antioxidants to counteract their effects. So, the present study investigated the potential protective effects of vitamin E (50 mg/kg BW), tomato paste (in terms of 9 mg lycopene/kg BW) and their combination versus atrazine-induced changes in *Clarias gariepinus*. This species were exposed to sublethal doses of atrazine of 1.7 and 3.4 mg/L for 15 and 30 days. Atrazine significantly ($P < 0.05$) induced free radicals in serum constituents. It caused a significant ($P < 0.05$) increase in serum glucose, mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH) and lipid peroxidation in liver. Addition of vitamin E and/or tomato paste to the diet of atrazine-treated fish improved the serum glucose, MCV, MCH and LPO levels in comparison to control fish. On the other hand, atrazine significantly ($P < 0.05$) led to decline in serum total protein, total lipid, blood hemoglobin, total erythrocytes count, packed cell volume, and mean corpuscular hemoglobin concentration. These atrazine-induced parameters were improved with the dietary supplemented tomato paste and/or vitamin E.

Keywords: *Clarias*, Atrazine, Tomato-past, vitamin-E haematology, biochemistry.

INTRODUCTION

The aquatic ecosystems have known to receive a wide spectrum of pollutants, which may be introduced to it directly or indirectly. Pesticides are one of these pollutants representing a serious environmental problem (Fischer-Scherl et al., 1991; Joseph and Raj, 2011; Khidir et al., 2001; Mekkawy et al., 1996; Okayi et al., 2013a; Okayi et al., 2013b). The major source of contamination

by these pesticides is the deposits resulting from their application to control agriculture pests and harmful aquatic herbs (Egaas et al., 1993; Fischer-Scherl et al., 1991; Mekkawy et al., 1996; Yadav et al., 2010). The pesticides have different diverse impacts on the aquatic animals especially fishes which are of economic importance and high value from the point of biological conservation. Different worldwide efforts and scientific studies have been carried out to evaluate and hence prevent the adverse impacts of these chemicals to the extent of their banning especially in the advanced countries (EUC, 2007).

*Corresponding Author E-mail: iaa_mekkawy@yahoo.co.uk

In Egypt, waste water and agriculture drains containing pesticides and heavy metals residue are discharged directly into the Nile and its tributaries and the northern lakes (Mekkawy et al., 1996). Pesticides enter these water bodies from industrial planes, aerial spraying of field and water drainage system (El-Elaimy et al., 1990). The increasing use of pesticides in insect and herbal control as well as in public and private gardens has resulted in a great danger to aquatic organisms and man (El-Missiry and Othman, 1993).

Atrazine (AT) is one of the herbicides used in Egypt. It belongs to the s-triazine family of herbicides, which are one of the most significant water pollutants in rain, fresh, marine and ground waters (Egaas et al., 1993; Felding, 1992; Fischer-Scherl et al., 1991; Tasli et al., 1996). Atrazine has been detected in various concentrations in rainwater, surface water, ground water and drinking water reservoirs (Braun et al., 1987; Haberer and Normann, 1988; Oehmichen and Haberer, 1986). Fish species are one of the aquatic organism which suffer severely from the herbicides especially atrazine. Different studies have been carried out to evaluate its adverse impacts on the biological system of fishes and to determine the corresponding biological, molecular and genetic biomarkers (e.g. (Khidr et al., 2001; Nwani et al., 2010; Okayi et al., 2013a; Ovie et al., 2007; Wassif et al., 2000; Yadav et al., 2010). Also, further studies are published to counteract the adverse impacts of pollutants especially in aquaculture through the enhancement of the natural antioxidant system and addition of different exogenous antioxidants in diets (Mekkawy et al., 2011; 2012).

Carotenoids are a family of fat-soluble pigments found in tomatoes and its products, some other fruits and vegetables. Many studies have investigated the potentials of these carotenoids in the oxidative stress. Tomato carotenoids include lycopene and other similar carotenoids (Cohen, 2002; Mekkawy et al., 2011; 2012; Tapiero et al., 2004; Visioli et al., 2003). Recently, tomato paste and its individual carotenoids have received particular attention in counteracting pollutant-induced oxidative stress especially in human and rats because of their highly efficient scavenging capacity for singlet-oxygen and free radicals (Heber and Lu, 2002; Mekkawy et al., 2011; 2012; Stahl and Sies, 2003; Wertz et al., 2004). The potentials of tomato and their products and carotenoids especially lycopene in treatment of different types of human cancers were evident especially in prostate cancers (Boileau et al., 2003; Mekkawy et al., 2011). Lycopene can be considered as a biomarker for the additive or synergistic anticancer effects of tomato phytonutrients. Boileau et al. (2003), supporting this observation, concluded that consumption of tomato powder but not lycopene inhibited prostate carcinogenesis. Mekkawy et al. (2011; 2012) referred to its validity in improving liver function and other biochemical and liver histological characteristics of cadmium-exposed *Oreochromis niloticus*.

Vitamin E is an indispensable nutrient required to maintain flesh quality, immunity, normal resistance of red blood corpuscles to haemolysis, permeability of capillaries and heart muscle (Halver, 2002). It functions as a lipid soluble antioxidant and protects biological membranes, lipoproteins and lipid stores against oxidation. Its main function is to protect unsaturated fatty acids against free radical-mediated oxidation (Hamre et al., 1998). Different studies have been recorded its antioxidant roles in preventing and improving pollutant adverse effects including those induced by atrazine on the biological systems of aquatic animals.

According to the aforementioned findings and low price of tomatoes in a definite season in Egypt, the present work was suggested and aimed to study the protective roles of tomato paste and/or vitamin E on atrazine-induced changes in some haematological and biochemical parameters of the Nile catfish, *Clarias gariepinus*.

MATERIALS AND METHODS

Sample collection and treatment manipulation

One hundred and twenty healthy fish of The Nile catfish, *Clarias gariepinus* were caught from the River Nile at Assiut, Egypt. Fishes immediately were transported to the fish laboratory in the Department of Zoology, Faculty of Science, Assiut University. The experimental fishes were reared in aerated glass tanks (100 L capacity) and acclimatized for two weeks before being used in the experimental study. The experimental fish fed pellets at a rate of 4.5 % of wet weight twice daily. Feces and residual food were aspirated regularly. The water temperature, pH and dissolved oxygen (DO) concentrations were measured daily ($18.44 \pm 1.43^{\circ}\text{C}$, 6.94 ± 0.11 pH, 6.71 ± 2.52 mg L⁻¹ DO) with photoperiod of 12 light/12 dark.

Preparation of tomato paste to adjust the lycopene dose

Preparation of tomato paste to adjust the lycopene dose follow Mekkawy et al. (2011). Tomatoes used for the experiment were obtained from the local market. Fresh peeled, deseeded tomatoes were pulped well to a smooth consistency in a Warring blender. The lycopene content in tomato paste was estimated spectrophotometrically according to the methods of Ranganna (1976) and Choudhari and Anantharayan (2007). The lycopene concentration in the tomato paste was 30.28 mg kg⁻¹. In addition to lycopene, tomato paste composition include water, proteins, carbohydrates, fibers, calcium, potassium, zinc, copper, manganese, iron, vitamin C, vitamin E, β -carotenoids and other

Table 1. The fish groups exposed to atrazine (AT1, AT2) doses, tomato paste (TP), vitamin E (VE) and their combinations.

Treatments	C	VE	TP	TP+ VE	AT1	AT1+ VE	AT1+ TP	AT1+T P+VE	AT2	AT2+ VE	AT2+ TP	AT2+ TP+ VE
Atrazine (mg/L)	0	0	0	0	1.7	1.7	1.7	1.7	3.4	3.4	3.4	3.4
Vitamin E (mg/kg)	0	50	0	50	0	50	0	50	0	50	0	50
Tomato paste (mg/kg)	0	0	9	9	0	0	9	9	0	0	9	9

C= control, VE= vitamin E, TP= tomato paste and AT1, AT2= atrazine doses

phytonutrients (Okajima et al., 1998).

Experimental design

Fishes were weighed, measured and classified randomly into 12 groups (10 fishes/tank) according to doses of atrazine (AT), tomato paste (TP), vitamin E (VE) and their combinations (Table 1). The diets were pelleted after addition of tomato paste and vitamin E doses and addition of suitable amounts of molasses and water. The diets were dried at room temperature and stored in small bags for fish feeding.

Stock solution of the herbicide, atrazine were prepared and stored in clean glass bottles and diluted to two sublethal concentrations of 1.7 and 3.4 mg L⁻¹ (96h- LC50 equal 7.6 mg L⁻¹ for catfish (WHO/FAO, 1996). Atrazine doses were prepared and added constantly to the aquarium for four weeks. The test water was replaced daily with the required amount of stock solution to prevent deterioration of water quality and replenish atrazine levels.

Blood analyses

After 15 and 30 day periods, blood samples of the control and treated fish (3 fish/treatment) were collected from caudal vein of the fish in a small plastic tubes containing heparin solution (0.2 mL mL⁻¹ blood) as anticoagulant. These blood samples were used for determining erythrocyte count according to Dacie and Lewis (1984) using hemocytometer. Haemoglobin (Hb) was estimated where it was converted into red cyanomethemoglobin under the influence of potassium ferricyanide and potassium cyanide according to Vankampen (1961). Haemoglobin level was determined by using suitable Kits (Diamond Diagnostics, Egypt) according to Stoskopf (1993). Haematocrit value (Hct) was calculated according to the formula mentioned by Stoskopf (1993). Mean corpuscular volume (MCV), Mean corpuscular

hemoglobin (MCH) and Mean corpuscular hemoglobin concentration (MCHC) were calculated according to Cloes (1980).

Some other blood samples were collected and left to coagulate for 15-20 minutes at 4°C prior to centrifugation for 20 minutes at 3000 rpm to separate serum. The fresh serum was subjected to biochemical analysis. Serum glucose (mg L⁻¹) was determined, using assay kit supplied by (Spectrum Diagnostics, Egypt). Total protein (g 100mL⁻¹) content and total lipids (g L⁻¹) content were determined colorimetrically using assay kit supplied by Diamond Diagnostics, Egypt. The samples were measured by spectrophotometer (Ultrospec 3100 pro, Biochrom Ltd).

Lipid peroxidation measurement

Lipid peroxidation was measured according to the method of Ohkawa *et al.* (1979). 10(w/v) tissue homogenate from liver was used (this homogenate contained 1% v/v dimethyl sulfoxide to prevent further oxidation). 0.2 ml aliquots of tissue homogenate were added to 0.2 ml sodium dodecyl sulfate solution (8.1% w/v) and 1.5 ml acetic acid solution (20% v/v). The mixture was made up to 4.0 ml with distilled water and heated to 95 °C for 1 h. The samples were cooled, centrifuged at 2000 rpm for 10 min. and measured at 532 nm using spectrophotometer (Ultrospec 3100 pro, Biochrom Ltd). The results were expressed as nmol malondialdehyde formation per g tissue.

Statistical analysis

The basic statistics, means, standard errors and ranges of the measured parameters were estimated. The patterns of variation due to atrazine, tomato paste and vitamin E doses and their combinations were studied by three and four-way analysis of variance using the SPSS transformed data. So, the homogeneity of variance was

Table 2. Values of blood constituent parameters of *Clarias gariepinus* exposed to atrazine, vitamin E, tomato paste and their combinations for 15 days.

Treatments Parameters	C	VE	TP	TP+VE	AT1	AT1+VE
RBCs(million/mm ³)	3.18±0.02 ^b (3.15-3.21)	3.41±0.24 ^d (3.00-3.82)	3.65±0.03 ^b (3.61-3.70)	2.23±0.05 ^a (2.13-2.32)	1.90±0.42 ^a (1.17-2.63)	2.27±0.23 ^a (1.86-2.67)
Hb(g/100ml blood)	8.32±0.47 ^{ab} (7.50-9.15)	10.61±0.20 ^c (10.26-10.96)	12.49±0.86 ^d (11.00-13.97)	10.79±0.18 ^c (10.52-11.15)	7.32±0.07 ^{ab} (7.20-7.43)	8.80±0.12 ^b (8.60-9.00)
Hct(%)	24.10±1.33 ^{ab} (21.80-26.39)	30.51±0.57 ^c (29.53-31.49)	35.77±2.40 ^d (31.60-39.92)	31.00±0.51 ^c (30.26-31.99)	21.29±0.18 ^{ab} (20.96-21.60)	25.44±0.32 ^b (24.88-26.00)
MCV(μ ³)	75.80±4.41 ^a (68.55-83.78)	90.44±7.10 ^a (77.30-101.69)	97.98±6.94 ^a (85.41-109.36)	139.52±5.77 ^a (130.41-150.20)	125.15±30.09 ^a (79.70-182.02)	115.83±12.58 ^a (93.18-136.77)
MCH(μμ ³)	26.17±1.58 ^a (23.58-29.02)	31.46±2.48 ^a (26.86-35.37)	34.21±2.47 ^a (29.73-38.27)	48.54±2.03 ^a (45.34-52.30)	43.02±0.34 ^a (27.38-62.56)	39.72±4.36 ^a (32.21-47.31)
MCHC(%)	34.52±0.07 ^{bc} (34.40-34.63)	34.78±0.02 ^{de} (34.75-34.81)	34.91±0.06 ^e (34.81-35.00)	34.79±0.01 ^{de} (34.77-34.82)	34.37±0.01 ^{abc} (34.35-34.39)	34.59±0.01 ^{cd} (34.57-34.62)
Gl(mg/dl)	25.08±1.75 ^a (22.05-30.11)	26.90±0.53 ^{ab} (25.98-27.82)	27.31±0.61 ^{ab} (26.25-30.36)	47.50±2.26 ^{cd} (43.25-50.96)	41.61±3.64 ^c (35.43-48.03)	38.58±2.88 ^{abc} (33.59-43.57)
TP(g/dl)	5.43±0.05 ^{de} (5.34-5.51)	6.41±0.01 ^{ef} (6.39-6.43)	5.13±0.16 ^{cd} (4.85-5.40)	6.62±0.25 ^f (6.18-7.06)	4.57±0.45 ^{bcd} (3.82-5.36)	4.27±0.11 ^{bcd} (4.07-4.46)
TL(g/l)	8.32±0.13 ^d (8.10-8.54)	8.37±0.27 ^d (7.89-8.84)	7.10±0.66 ^{cd} (5.95-8.24)	8.05±0.25 ^d (7.61-8.49)	7.08±0.07 ^{cd} (6.98-7.21)	8.19±0.12 ^d (7.98-8.40)
LPO(nmol/mg tissue)	1.39±0.59 ^{ab} (0.37-2.41)	1.13±0.11 ^a (0.94-1.32)	1.49±0.19 ^{ab} (1.15-1.82)	1.5±0.09 ^{ab} (1.32-1.61)	4.20±0.33 ^d (3.63-4.77)	2.61±0.18 ^{bc} (2.24-2.79)

The data are presented as Means ± S.E. (Min-Max).

Different letters indicate significance at p<0.05

Red blood cells (RBC), Haemoglobin (Hb), Hematocrit (Hct), Mean corpuscular volume (MCV), Mean corpuscular haemoglobin (MCH), Mean corpuscular haemoglobin concentration (MCHC), Glucose (Gl), Total protein (TP), Total lipids (TL) and Lipid peroxidation (LPO).

Table 2. Cont.

Treatments Parameters	AT1+TP	AT1+TP+VE	AT2	AT2+VE	AT2+TP	AT2+TP+VE
RBCs(million/mm ³)	2.17±0.21 ^a (1.80-2.54)	1.97±0.25 ^a (1.53-2.40)	1.49±0.21 ^a (1.13-1.85)	1.98±0.12 ^a (1.76-2.19)	1.71±0.03 ^a (1.66-1.75)	1.73±0.01 ^a (1.70-1.75)
Hb(g/100ml blood)	7.68±0.22 ^{ab} (7.30-8.05)	7.00±0.18 ^a (6.69-7.30)	6.84±0.21 ^{ab} (6.47-7.20)	7.30±0.40 ^{ab} (6.58-7.98)	7.49±0.30 ^{ab} (7.02-8.00)	6.90±0.46 ^a (6.10-7.70)
Hct(%)	22.29±0.61 ^{ab} (21.24-23.34)	20.39±0.49 ^a (19.53-21.24)	19.94±0.59 ^{ab} (18.92-20.96)	21.18±1.13 ^{ab} (19.22-23.15)	21.77±0.79 ^{ab} (20.46-23.20)	20.12±1.29 ^a (17.88-22.36)
MCV(μ ³)	104.56±9.83 ^a (91.89-123.91)	106.57±11.40 ^a (88.50-127.66)	139.83±21.46 ^a (102.25-176.57)	107.30±1.04 ^a (105.68-109.23)	127.68±5.60 ^a (116.89-135.67)	116.42±6.52 ^a (105.18-127.77)
MCH(μμ ³)	36.00±3.38 ^a (31.69-42.67)	36.56±3.88 ^a (30.42-43.73)	47.94±7.38 ^a (34.97-60.53)	36.87±0.30 ^a (36.44-37.39)	43.92±1.98 ^a (40.11-46.78)	39.92±2.34 ^a (35.88-44.00)
MCHC(%)	34.43±0.03 ^{abc} (34.37-34.49)	34.31±0.03 ^{ab} (34.25-34.37)	34.30±0.04 ^{abc} (34.20-34.35)	34.36±0.07 ^{ab} (34.23-34.48)	34.40±0.05 ^{abc} (34.32-34.48)	34.30±0.09 ^a (34.12-34.44)
Gl(mg/dl)	46.98±5.46 ^{cd} (38.58-57.22)	64.29±2.26 ^e (60.52-68.32)	58.85±1.93 ^c (55.40-62.08)	41.21±0.61 ^c (40.16-42.26)	39.77±0.53 ^{bc} (38.85-40.68)	65.11±6.53 ^e (55.38-77.53)
TP(g/dl)	4.53±0.16 ^{cd} (4.26-4.82)	4.35±0.07 ^{bcd} (4.23-4.46)	4.11±0.54 ^{cd} (3.17-5.04)	3.00±0.33 ^a (2.42-3.58)	3.93±0.33 ^{abc} (3.35-4.50)	3.77±0.15 ^{ab} (3.52-4.01)
TL(g/l)	4.59±0.06 ^a (4.47-4.68)	5.51±0.25 ^{ab} (5.07-5.95)	6.06±0.06 ^{cd} (5.95-6.16)	5.27±0.36 ^{ab} (4.65-5.88)	5.85±0.29 ^b (5.35-6.34)	4.55±0.15 ^a (4.30-4.79)
LPO(nmol/mg tissue)	6.47±0.21 ^e (6.12-6.83)	2.64±0.58 ^{bc} (1.99-3.80)	3.23±0.18 ^{cd} (2.92-3.55)	1.21±0.04 ^a (1.15-1.30)	5.70±0.10 ^e (5.53-5.85)	2.1±0.23 ^{abc} (1.70-2.50)

The data are presented as Means ± S.E. (Min-Max).

Different letters indicate significance at p<0.05

Red blood cells (RBCs), Haemoglobin (Hb), Hematocrit (Hct), Mean corpuscular volume (MCV), Mean corpuscular haemoglobin (MCH), Mean corpuscular haemoglobin concentration (MCHC), Glucose (Gl), Total protein (TP), Total lipids (TL) and Lipid peroxidation (LPO).

Table 3. Values of blood constituent parameters of *Clarias gariepinus* exposed to atrazine, vitamin E, tomato paste and their combinations for 30 days.

Treatments Parameters	C	VE	TP	TP+VE	AT1	AT1+VE
RBCs(million/mm ³)	3.40±0.16 ^d (3.12-3.67)	3.31±0.03 ^d (3.25-3.36)	3.77±0.15 ^d (3.52-4.01)	2.23±0.10 ^{abc} (2.06-2.39)	1.92±0.11 ^{abc} (1.75-2.12)	2.55±0.12 ^c (2.35-2.75)
Hb(g/100ml blood)	9.11±0.34 ^d (8.53-9.70)	9.32±0.31 ^d (8.79-9.85)	12.15±0.68 ^e (10.96-13.33)	8.42±0.38 ^{bcd} (7.76-9.08)	6.66±0.21 ^a (6.29-7.02)	7.41±0.22 ^{abc} (7.02-7.79)
Hct(%)	26.31±0.95 ^d (24.68-27.96)	26.90±0.86 ^d (25.41-30.38)	34.81±1.91 ^e (31.49-38.12)	24.38±1.07 ^{bcd} (22.53-26.22)	19.44±0.59 ^a (18.41-20.46)	21.54±0.62 ^{abc} (20.46-22.61)
MCV(μ ³)	77.53±0.85 ^a (76.19-79.12)	81.30±1.81 ^a (78.19-84.46)	92.29±1.62 ^{ab} (89.45-95.07)	109.46±0.13 ^{ab} (109.31-109.72)	101.35±2.57 ^{ab} (96.49-105.21)	84.92±5.34 ^a (74.39-91.69)
MCH(μμ ³)	26.84±0.27 ^a (26.43-27.34)	30.18±0.66 ^a (27.05-29.32)	32.20±0.61 ^{ab} (31.15-33.24)	37.81±0.10 ^{ab} (37.67-37.99)	34.70±0.84 ^{ab} (33.11-35.94)	29.20±1.86 ^a (25.53-31.53)
MCHC(%)	34.63±0.04 ^d (34.56-34.69)	34.65±0.03 ^d (34.59-34.71)	34.89±0.04 ^e (34.81-34.96)	34.54±0.05 ^{cd} (34.45-34.62)	34.24±0.05 ^a (34.16-34.32)	34.39±0.04 ^{abc} (34.32-34.45)
Gl(mg/dl)	26.59±1.05 ^a (24.77-30.40)	30.43±1.17 ^a (30.40-32.46)	27.95±0.96 ^a (26.30-29.61)	26.15±0.44 ^a (25.38-26.89)	58.76±5.67 ^c (48.94-68.58)	37.95±1.34 ^{ab} (35.45-40.02)
TP(g/dl)	6.06±0.13 ^e (5.82-6.26)	7.62±0.43 ^f (6.88-8.36)	7.26±0.40 ^f (6.56-7.96)	5.08±0.07 ^{cde} (5.00-5.21)	4.25±0.15 ^{bc} (4.02-4.50)	5.30±0.20 ^{cde} (4.95-5.65)
TL(g/l)	8.53±0.25 ^d (8.12-8.97)	9.27±0.44 ^d (8.50-10.04)	8.65±0.04 ^d (8.58-8.72)	9.23±0.25 ^d (8.78-9.63)	6.07±0.12 ^{abc} (5.85-6.30)	5.03±0.04 ^{ab} (4.96-5.09)
LPO(nmol/mg tissue)	2.77±0.06 ^a (2.66-2.88)	6.83±0.05 ^{bcd} (6.75-6.92)	6.30±0.44 ^{bc} (5.40-6.79)	5.04±0.59 ^b (4.01-6.07)	6.73±0.35 ^{bcd} (6.12-7.34)	7.71±0.49 ^{bc} (7.00-8.64)

The data are presented as Means ± S.E. (Min-Max).

Different letters indicate significance at p<0.05

Red blood cells (RBCs), Haemoglobin (Hb), Hematocrit (Hct), Mean corpuscular volume (MCV), Mean corpuscular haemoglobin (MCH), Mean corpuscular haemoglobin concentration (MCHC), Glucose (Gl), Total protein (TP), Total lipids (TL) and Lipid peroxidation (LPO).

Table 3. Cont.

Treatments Parameters	AT1+TP	AT1+TP+VE	AT2	AT2+VE	AT2+TP	AT2+TP+VE
RBCs(million/mm ³)	2.49±0.02 ^{bc} (2.45-2.53)	1.90±0.36 ^{abc} (1.27-2.53)	1.44±0.01 ^a (1.42-1.45)	2.00±0.12 ^{abc} (1.80-2.20)	2.42±0.16 ^{bc} (2.15-2.70)	1.70±0.29 ^{ab} (1.19-2.20)
Hb(g/100ml blood)	8.91±0.20 ^d (8.56-9.26)	8.10±0.13 ^{bcd} (7.86-8.31)	6.65±0.10 ^a (6.47-6.82)	7.16±0.08 ^{ab} (7.02-7.30)	8.63±0.23 ^{cd} (8.24-9.05)	8.11±0.31 ^{bcd} (7.56-8.64)
Hct(%)	25.75±0.57 ^d (24.77-26.73)	23.49±0.37 ^{bcd} (22.81-24.07)	19.41±0.30 ^a (18.92-19.90)	20.85±0.23 ^{ab} (20.46-21.24)	24.96±0.66 ^{cd} (23.87-26.15)	23.52±0.87 ^{bcd} (21.97-24.99)
MCV(μ ³)	103.38±1.31 ^{ab} (101.09-105.64)	133.17±25.15 ^{ab} (93.25-179.59)	135.13±2.37 ^{ab} (130.46-138.17)	104.81±4.94 ^{ab} (96.55-113.64)	103.90±6.20 ^{ab} (96.81-116.26)	158.83±31.25 ^b (107.24-210.02)
MCH(μμ ³)	35.77±0.48 ^{ab} (34.94-36.60)	45.93±8.65 ^{ab} (32.17-61.89)	46.27±0.84 ^{ab} (44.62-47.36)	35.99±1.68 ^{ab} (33.18-39.00)	35.92±2.15 ^{ab} (33.52-40.19)	51.36±10.84 ^b (37.00-72.61)
MCHC(%)	34.60±0.03 ^d (34.56-34.65)	34.50±0.02 ^{bcd} (34.46-34.53)	34.24±0.02 ^a (34.20-34.30)	34.34±0.01 ^{ab} (34.32-34.37)	34.57±0.03 ^d (34.52-34.62)	34.49±0.05 ^{bcd} (34.41-34.57)
Gl(mg/dl)	30.25±0.79 ^a (26.89-29.61)	38.82±0.79 ^{ab} (37.46-40.18)	80.63±5.72 ^e (69.42-88.22)	34.75±2.27 ^a (30.82-38.67)	29.46±2.36 ^a (25.38-33.54)	47.96±1.41 ^b (45.36-50.21)
TP(g/dl)	5.65±0.19 ^{de} (5.31-5.98)	4.58±0.23 ^{bcd} (4.21-5.00)	2.80±0.18 ^a (2.48-3.11)	4.86±0.09 ^{cd} (4.68-4.97)	3.68±0.01 ^{ab} (3.66-3.69)	3.69±0.31 ^{ab} (3.13-4.20)
TL(g/l)	8.42±0.42 ^d (7.69-9.15)	7.20±0.19 ^c (6.84-7.50)	5.32±0.33 ^{ab} (4.74-5.89)	4.78±0.44 ^a (4.06-5.59)	5.13±0.05 ^{ab} (5.04-5.21)	6.26±0.41 ^{bc} (5.55-6.96)
LPO(nmol/mg tissue)	8.05±0.29 ^{cd} (7.55-8.56)	12.58±0.81 ^f (11.17-13.99)	8.43±0.22 ^d (8.05-8.81)	15.37±0.17 ^g (15.07-15.66)	10.63±0.42 ^e (9.82-11.25)	8.27±0.02 ^d (8.23-8.31)

The data are presented as Means ± S.E. (Min-Max).

Different letters indicate significance at p<0.05

Red blood cells (RBCs), Haemoglobin (Hb), Hematocrit (Hct), Mean corpuscular volume (MCV), Mean corpuscular haemoglobin (MCH), Mean corpuscular haemoglobin concentration (MCHC), Glucose (Gl), Total protein (TP), Total lipids (TL) and Lipid peroxidation (LPO).

package (SPSS, 1998) at the 0.05 significance level. Null hypothesis for raw, log-transformed and SQRT- Levene's test of equality of error variance of the dependent variables was applied, with rejection of the assumed for raw data. The pattern of variations was also recorded by one-way analysis of variance, revealing significant difference due to atrazine, tomato paste and vitamin E ($P < 0.0001$); The Tukey-HSD test was considered for multiple comparisons. Moreover, the Dunnett-T test was applied, measuring the control against other treatments.

RESULTS

Behavioral changes

After exposure to atrazine some of the fishes exhibited hyperactivity characterized by rapid and erratic swimming or darting, partial loss of equilibrium, rapid pectoral fins and opercular movements, reduction in the feeding activity, fins haemorrhage and loss of some skin parts especially in those exposed to atrazine without vitamin E and/or lycopene. Quick ventilation movements were recorded especially in aquaria with high concentration of atrazine. The mortality rate of 2.3 was recorded at the first period in aquaria with only atrazine. The dead fishes exhibited changes and abnormality in eyes, gills, gall bladder, spleen and liver color. On the other hand, fish groups exposed to the same doses of atrazine in combination with tomato paste and/or vitamin E did not show such behavioral abnormalities. Moreover, Atrazine-free fishes treated with vitamin E and/or tomato paste were noticed in better conditions.

Blood constituents

The basic data are given in Tables 2 and 3. Correlation matrix of blood constituents parameters in the first and second periods of exposure are given in Tables 4 and 5. Many significant correlations between most of such parameters were recorded. For example, RBCs were found to be correlated with all parameters.

A- Haematological parameters

The normal values of red-blood-cells (RBCs) count, haemoglobin (Hb) content and haematocrit (Hct) value of *Clarias gariepinus* in the two periods are given in Tables 2 and 3. The atrazine main effect, was highly significant ($P < 0.0001$) in both periods for the three previous parameters. The tomato paste main effect was significant in both periods except for RBCs. Vitamin E main effect was significant only at the second period for RBCs, Hb and Hct. TP-VE interaction effect was highly significant in both periods. AT-TP-VE interaction was only significant in

the second period. The time of exposure main effect was not significant and positive correlations were recorded between RBCs, Hb and Hct in the two periods (r RBCs= 0.948, r Hb and r Hct= 0.763).

The normal value of mean corpuscular value, mean corpuscular haemoglobin and mean corpuscular haemoglobin concentration of *Clarias gariepinus* in the two periods are given in Tables 2 and 3. The atrazine main effect was highly significant ($P < 0.0001$) in the two periods for MCHC while MCV and MCH showed significant effects only in the second period. The tomato paste main effect was significant for MCH, MCV and MCHC in the second period only. Vitamin E had no significant effect at the two periods except for MCHC. TP-VE interaction effect was significant for MCH, MCV and MCHC in the second period while MCHC showed significant effect in the first period only. AT-TP-VE interaction didn't show significant effect for the three parameters. The time of exposure main effect was not significant and positive correlations were recorded between MCV, MCH and MCHC in the two periods (r MCV= 0.544, r MCH= 0.517 and r MCHC= 0.689). Supplementation of tomato paste and/or vitamin E improved the haematological parameters of AT- exposed fishes (Tables 2 and 3).

B- Biochemical parameters

Glucose level

The normal glucose level of *Clarias gariepinus* in the two periods is given in Tables 2 and 3. Atrazine main effect was significantly increased in both periods ($P < 0.0001$). The main effect of tomato paste and/or vitamin E in the two periods was highly significant ($P < 0.0001$). AT-TP-VE interaction showed significant effect in the two periods. The time of exposure main effect was significant ($P < 0.0001$) and a positive correlation was obtained between glucose levels in the two periods ($r = 0.526$). Dietary supplementation with vitamin E and/or tomato paste improved AT-induced hyperglycemia in comparison with control.

Total protein level

The total protein level of *Clarias gariepinus* in the two periods is given in Tables 2 and 3. Atrazine reflected high significant decrease in total protein level at both periods ($P < 0.0001$). The main effect of tomato paste or vitamin E in the two periods was not significant. Highly significant interaction effect between AT- TP- VE was recorded in the second period only. The time of exposure main effect was highly significant ($P < 0.0001$) and a positive correlation was obtained between the total protein levels total protein in AT-exposed fish.

Table 4. Correlation matrix of blood constituent parameters of *Clarias gariepinus* subjected to different treatments of atrazine, tomato paste, vitamin E and their combinations for 15 days.

Parameters	RBCs	HB	HCT	MCV	MCH	MCHC	Glucose	TP	TL	LPO
RBCs	1.000									
HB	0.721**	1.000								
HCT	0.721**	1.000**	1.000							
MCV	-0.730**	-0.113	-0.113	1.000						
MCH	-0.711**	-0.086	-0.086	1.000**	1.000					
MCHC	0.725**	0.980**	0.980**	-0.127	-0.100	1.000				
Glucose	-0.629**	-0.577**	-0.577**	0.256	0.241	-0.630**	1.000			
TP	0.573**	0.656**	0.656**	-0.165	-0.156	0.685**	-0.357*	1.000		
TL	0.524**	0.591**	0.591**	-0.121	-0.104	0.635**	-0.620**	0.686**	1.000	
LPO	-0.435**	-0.421*	-0.421*	0.215	0.205	-0.391*	0.206	-0.277	-0.430**	1.000

* Significant at P<0.05

** Significant at P<0.01

Table 5. Correlation matrix of blood constituent parameters of *Clarias gariepinus* subjected to different treatments of atrazine, tomato paste, vitamin E and their combinations for 30 days.

Parameters	RBCs	HB	HCT	MCV	MCH	MCHC	Glucose	TP	TL	LPO
RBCs	1.000									
HB	0.789**	1.000								
HCT	0.789**	1.000**	1.000							
MCV	-0.758**	-0.254	-0.254	1.000						
MCH	-0.746**	-0.235	-0.235	1.000**	1.000					
MCHC	0.780**	0.972**	0.972**	-0.257	-0.238	1.000				
Glucose	-0.608**	-0.586**	-0.586**	0.399*	0.386*	-0.681**	1.000			
TP	0.827**	0.707**	0.707**	-0.538**	-0.527**	0.701**	-0.646**	1.000		
TL	0.566**	0.627**	0.627**	-0.240	-0.230	0.674**	-0.495**	0.681**	1.000	
LPO	-0.485**	-0.320	-0.320	0.362*	0.358*	-0.320	0.125	-0.359*	0.575**	1.000

* Significant at P<0.05

** Significant at P<0.01

in the two periods ($r = 0.608$). In the present study, diet supplementation with tomato paste and/or vitamin E for 15 and 30 days showed a marked elevation in the level of

Total lipids level

The total lipids level of *Clarias gariepinus* in the two periods is given in Tables 2 and 3. Atrazine reflected high significant decrease in total lipids level ($P < 0.0001$) in the two periods of exposure. The main effect of tomato paste in the two periods was significant ($P < 0.0001$). Vitamin E and its interaction with tomato paste didn't show significant effect in the two periods. Highly significant interaction effect between AT-TP and AT-VE was recorded in the two periods. The time of exposure main effect was highly significant ($P < 0.0001$) and a positive correlation was obtained between the total lipids levels in the two periods ($r = 0.387$). Dietary supplementation of tomato paste and vitamin E corrected markedly the level of total lipids in comparison of that of control and AT-

treated group ($P < 0.0001$) especially in the second period.

Lipid peroxidation measurement

The liver lipid peroxidation (LPO) values are given in Tables 2 and 3. The main effects of AT, TP, VE and their interactions were highly significant ($P < 0.0001$) in the two periods except for AT-VE interaction in the second period. The time of exposure main effect was highly significant ($P < 0.0001$) and a positive correlation was recorded between LPO in the two periods ($r = 0.17$). The values of lipid peroxidation were significantly ($P < 0.0001$) decreased in liver of AT-exposed fishes fed diets supplemented with vitamin E and/or tomato paste especially in the first period.

DISCUSSION

In the present study, *Clarias gariepinus* exhibited some

behavioral changes due to exposure to sublethal concentrations of atrazine. Similar behavioral changes were recorded by El-Banhawy et al. (1986); Abdel-Rahman (1997); Wassif et al. (2000); Hussein and Mekkawy (2001); Saha and Kaviraj (2003); Viran et al. (2003); Calta and Ural (2004); Köprücü et al. (2006), Sobha et al. (2007) and Mekkawy et al. (2011) in fishes exposed to pesticides and heavy metals. Such atrazine-induced behavioral changes were counteracted by dietary supplementation with tomato paste and/or vitamin E; a result confirmed by Mekkawy et al. (2011) for cadmium induced behavioral changes of *O. niloticus*. Moreover, the control fishes of *C. gariepinus* and *O. niloticus* treated with vitamin E and/or tomato paste were noticed in better conditions.

The atrazine-induced decrease in RBCs, Hb and Hct of *Clarias gariepinus* recorded in the present study is in agreement with those of Mekkawy et al. (1996) working on *O. niloticus* and *Chrysichthys auratus* exposed to atrazine, Köprücü et al. (2006) working on *Silurus glanis* exposed to diazinon and Siang et al. (2007) working on *Monopterus albus* exposed to endosulfan. The reduction of these parameters at sublethal levels of these pesticides might be due to the destruction of mature RBCs and the inhibition of erythrocyte production due to reduction of haem synthesis that is affected by these pollutants (Khadre, 1988; Wintrobe, 1978). Also, this reduction may be indirectly attributed to pesticide-induced decrease in growth and other food utilization parameters which results in severe anemia (James and Sampath, 1999). Heath (1987) and Abo-Hegab et al. (1993) interpreted stress-induced decrease in the haemoglobin and haematocrit values in terms of haem dilution of blood and elimination of RBCs as well as disequilibrium of the osmotic pressure inside and outside the blood cell.

The atrazine-induced changes towards increase or decrease in MCV, MCH and MCHC values were reported in the present work. Similar findings were recorded by (Mekkawy et al., 1996) working on *Oreochromis niloticus* and *Chrysichthys auratus* and (Köprücü et al., 2006) working on *Silurus glanis* after exposure to the pesticides, atrazine and diazinon respectively.

Adeyemo (2007) found significant increase in MCV, MCH and MCHC values of *Clarias gariepinus* exposed to lead while Siang et al. (2007) working on *Monopterus albus* exposed to endosulfan found decrease in MCH and MCHC and increase in MCV values.

These chemicals-induced alterations in MCV, MCH and MCHC were attributed to direct or feedback responses of structural damage to RBC membranes resulting in haemolysis and impairment in hemoglobin synthesis, stress-related release of RBCs from the spleen and hypoxia (Marie et al., 1998; Shah, 2006). Except for those of Mekkawy et al. (2012) on *O. niloticus*, no literature was in hand with regards to lycopene and tomato paste benefits for fish stressed haematological parameters. However, many studies were in concern with

the beneficial supplementation of vitamin E to counteract the pesticide sublethal effects on fish haematology (Andrade et al., 2007; Montero et al., 2001) confirming the present results. In the present study, atrazine-induced hyperglycemia was revealed in *Clarias gariepinus*. Similar findings were observed by Mekkawy et al. (1996); Elowa and El-Elaimy (2001); Fayed et al. (2001); Das and Mukherjee (2003) and Fouda (2004) after exposure to different doses of pesticides. The source of such hyperglycemia seems to be due to the liver glycogenolysis, resulting from the increased plasma catecholamines and corticosteroid hormones (Mazeaud et al., 1977; Pickering, 1981) as well as amino acids through the activation of gluconeogenesis process (Abo-Hegab and Hanke, 1984). Tomato paste and/or vitamin E reduced atrazine-induced hyperglycemia revealed in *Clarias gariepinus*. Similar results were recorded by Mekkawy et al. (2011) for *O. niloticus* stressed by cadmium. Martins et al. (2007) reported a reduced blood glucose level after stress in Atlantic halibut (*Hippoglossus hippoglossus* L.) fed vitamin E supplemented diets.

In the present study, decrease in serum total protein level (hypoproteinemia) was recorded in atrazine-exposed *C. gariepinus*. Similar findings were recorded by Assem et al. (1992), Mekkawy et al. (1996; 2011), Das and Mukherjee (2003), Fouda (2004), Gad (2005) and Shalaby (2007) after exposure to different doses of pesticides and heavy metals. Such hypoproteinemia may be due to direct effect of the utilization of body protein as an energy supply to meet the increasing physiological demands to overcome the stress in the polluted medium (Abd El-Salam et al., 1994; El-Sayed et al., 1996; Fontana et al., 1998) hypoproteinemia may also be attributed to several pathological processes including plasma dissolution, renal damage and elimination in the urine, decreased liver protein synthesis, alteration in hepatic blood flow and / or hemorrhage into the peritoneal cavity and intestine (Keith and Weber, 1979). The present results and those of Mekkawy et al. (2011) confirm the validity of tomato paste and/or vitamin E in counteracting stress-induced hypoproteinemia. Elkomy and Hassan (2005) observed an increase in the stressed total protein level in male rat fed Tomato-juice as supplemented diets. Kalender et al. (2005) and Ogur et al. (2005) recorded an increase in total protein level after Diazinon-induced and Nitrate-induced stress in male rats fed vitamin E as supplemented diets.

In the present study, decrease in serum total lipids level was recorded in atrazine-treated *C. gariepinus*. Similar findings were observed by Mekkawy et al. (1996; 2011); Fayed et al. (2001) and Fouda (2004) after exposure to different doses of pesticides and heavy metals. Such induced decrease in serum total lipids level may be due to the increase in secretion of catecholamines and corticosteroids with enhanced metabolic rate and in turn reduced metabolic reserves (Fayed et al., 2001). Tomato paste and/or E were found

to be valid in counteracting stress-induced changes in total lipid level in the present work and that of Mekkawy et al. (2011). Yilmaz et al. (1997) found that vitamin E as a diet supplementation improved total lipids level in muscle and liver tissues of rats.

The present results have clearly demonstrated the ability of atrazine sublethal doses to induce oxidative stress in fish liver as evidenced by increased thiobarbituric acid reactive substance after atrazine-exposure. The rise in lipid peroxidation level in liver means a modification in the physical characteristics of cell membrane (Ursini et al., 1991) since lipid peroxidation leads to hydrolysis of phospholipids into hydroperoxy fatty acids (Salgo et al., 1993). The atrazine-induced oxidative stress in tissues by increasing lipid peroxidation and by altering the antioxidant status was postulated by many authors (Geret et al., 2002; Mekkawy et al., 2011; 2012; Nwani et al., 2010; Romeo et al., 2000; Romeo and Gnassia-Baralli, 1997; Sarkar et al., 1997).

In the present study, the level of lipid peroxidation was decreased in liver of atrazine-exposed fishes fed diets supplemented with vitamin E and/or tomato paste especially in the first period. Similar results were recorded by Mekkawy et al. (2011) in concern with cadmium-exposed *O. niloticus*. The benefits of vitamin E in elimination of stress-induced lipid peroxidation were recorded for different fish species including *Clarias gariepinus* (Baker and Davies, 1996; 1997), *Salmo salar* (Scaife et al., 2000), *Salmo gairdneri* (Chaiyapechara et al., 2003; Frigg et al., 1990; Hung and Slinger, 1982), hybrid tilapia, *Oreochromis niloticus* x *O. aureus* (Huang et al., 2003; 2004) and red hybrid tilapia, *Oreochromis* sp. (Wang et al., 2006). In rats, tomato and its products including lycopene were found by many authors to improve stress-induced lipid peroxidation (Atessahin et al., 2006; Bhuvaneshwaria et al., 2001; El-Demerdash et al., 2004; Elkomy and Hassan, 2005; Moreira et al., 2005; Velmurugan et al., 2001; 2002; Yilmaz et al., 2006).

REFERENCES

- Abd El-Salam MM, Mohamed GA, Lotfy SA (1994). The effect of deltamethrin on the levels of certain amino acids and some enzymes in catfish heart and serum. *Journal of Egyptian-German Society of Zoology*. 13, 345-363.
- Abdel-Rahman MA (1997). Toxicological studies of heavy metals on *Siganus rivulatus*., Vol. M.Sc. Alexandria University, Alexandria, Egypt.
- Abo-Hegab S, Hanke W (1984). The significance of cortisol for osmoregulation in carp (*Cyprinus carpio*) and tilapia (*Sarotherodon mossambicus*). *General and Comparative Endocrinology*. 54, 409-417.
- Abo-Hegab S, Kamel MG, Labib WD (1993). Some physiological and biochemical indices of the pesticide tamaron and bayluseide in fresh water fish *Tilapia Oreochromis niloticus*. *Proceedings of the Zoological Society, A.R.E.* 24, 183-197.
- Adeyemo OK (2007). Haematological Profile of *Clarias gariepinus* (Burchell, 1822) exposed to Lead. *Turkish J. Fisheries and Aquatic Sci.* 7, 163-169.
- Andrade JIA, Ono EA, Menezes GC, Brasil EM, Roubach R, Urbinati EC, Tavares-Dias M, Marcon JL, Affonso EG (2007). Influence of diets supplemented with vitamins C and E on pirarucu (*Arapaima gigas*) blood parameters. *Comparative Biochemistry and Physiology*. 146, 576-580.
- Assem H, Abo-Hegab S, Belal I (1992). Comparison of haematological effects of some toxicants on *Clarias gariepinus*. *Journal of Egyptian-German Society of Zoology*. 9, 33-50.
- Atessahin A, Karahan I, Türk G, Gür S, Yilmaz S, Çeribas IAO (2006). Protective role of lycopene on cisplatin-induced changes in sperm characteristics, testicular damage and oxidative stress in rats. *Reproductive Toxicology*. 21, 42-47.
- Baker RTM, Davies SJ (1996). Changes in tissue α -tocopherol status and degree of lipid peroxidation with varying α -tocopheryl acetate inclusion in diets for African catfish. *Aquacult. Nutr.* 2, 71-79.
- Baker RTM, Davies SJ (1997). The quantitative requirement for α -tocopherol for juvenile African catfish, *Clarias gariepinus* Burchell. *Animal Science (East Lothian, Scotland)*. 65, 135-142.
- Bhuvaneshwaria V, Velmurugana B, Balasenthila S, Ramachandranb CR, Naginia S (2001). Chemopreventive efficacy of lycopene on 7, 12-dimethylbenz [a] anthracene-induced hamster buccal pouch carcinogenesis. *Fitoterapia*. 72, 865-874.
- Boileau TWM, Liao ZKSSL Jr., EJW, SK, C (2003). Prostate carcinogenesis in N-methyl-N-nitrosourea (NMU)-testosterone treated rats fed tomato powder, lycopene, or energy-restricted diets. *J. National Cancer Institute (Bethesda, MD)*. 95, 1578-1586.
- Braun F, Schüssler W, Wehrle R (1987). Organische schadstoffe polychlorbiphenyle (PCB) und pestizide im kreislauf des wassers. *Bilanzierung und Bewertung. The Bavarian Water Research Institute, Munich*. 18-27.
- Calta M, Ural MS (2004). Acute toxicity of the synthetic pyrethroid deltamethrin to young mirror carp, *Cyprinus carpio*. *Fresenius Environmental Bulletin*. 13, 1179-1183.
- Chaiyapechara S, Casten MT, Hardy RW, Dong FM (2003). Fish performance, fillet characteristics, and health assessment index of rainbow trout (*Oncorhynchus mykiss*) fed diets containing adequate and high concentrations of lipid and vitamin E. *Aquaculture*. 219, 715-738.
- Choudhari SM, Ananthanarayan L (2007). Enzyme aided extraction of lycopene from tomato tissues. *Food Chemistry*. 102, 77-81.
- Cloes EH (1980). *Verterinary clinical pathology*. W. B. Saunder Company, London.
- Cohen LA (2002). A review of animal model studies of tomato communities in experimental ponds to atrazine, the most heavily used pesticide in the United States. *Ecology* 1982. 63, 1285-1293.
- Dacie JV, Lewis SM (1984). *Practical Haematology*. Churchill Living Stone, London.
- Das BK, Mukherjee S (2003). Toxicity of cypermethrin in *Labeo rohita* fingerlings: biochemical, enzymatic and haematological consequences. *Comparative Biochemistry and Physiology*. 134, 109-121.
- Egaas E, Skaare JU, Svendsen NO, Sandvik M, Falls JG, Dauterman WC, Collier TK, Netland J (1993). A comparative study of effects of atrazine on xenobiotic metabolizing enzymes in fish and insect, and of the in vitro phase II atrazine metabolism in some fish, insects, mammals and one plant species. *Comp Biochem Physiol C*. 106, 141-9.
- El-Banhawy MA, Al-Zahaby A, Shalaby A (1986). Histopathological studies on the effect of cyolane on the ileum of *Clarias lazira*. *Egyptian J. Histol.* 9, 77-85.
- El-Demerdash FM, Yousef MI, Kedwany FS, Baghdadi HH (2004). Cadmium-induced changes in lipid peroxidation, blood hematology, biochemical parameters and semen quality of male rats: protective role of vitamin E and β -carotene. *Food and Chemical Toxicology*. 42, 1563-1571.
- El-Elaimy IA, El-dib MA, Elowa SE (1990). Influence of solvents on the inhibitory potency of chloropyrifos and lannat on acetylcholinesterase from tissues of fresh water fish. *J. Egypt. Soc. Toxicol.* 15, 67-74.
- Elkomy MM, Hassan HA (2005). The role of tomato-juice as a protective agent against thioacetamide hepatotoxicity in male rats. *Comparative Physiology*. 46, 217-234.
- El-Missiry MA, Othman AI (1993). Influence of lannat on biochemical

- and haematological parameters in old rated. *J. Egyptian-German Soc. Zool. A*, 219-229.
- Elowa SHE, El-Elaimy IA (2001). Disturbance of some metabolic activities in liver of the freshwater fish, *Oreochromis niloticus* (Trewavas, 1983) under stress of pesticide-water pollution. *J. Egyptian-German Soc. of Zool.* 35, 23-39.
- El-Sayed AFM, El-Gobashy A, Al-Amoudi M (1996). Effects of pond depth and water temperature on the growth, mortality and body composition of Nile tilapia, *Oreochromis niloticus* (L.). *Aquaculture Research*. 27, 681-687.
- EUC, EU policy for a sustainable use of pesticides: the story behind the strategy. Office for Official Publications of the European Communities, Luxembourg (2007). pp. 26.
- Fayed HM, Zaghoul KH, Abd El-Monem S, Mohamed HA (2001). Biological responses of the Nile Tilapia, *Oreochromis niloticus* and the Nile catfish, *Clarias gariepinus* induced by agricultural and industrial pollutants. *J. Union of Arab Biol.* 16, 543-568.
- Felding G (1992). Leaching of atrazine into groundwater. *Pesticide Science*. 35, 39-41.
- Fischer-Scherl T, Veeseer A, Hoffmann RW, Kuhnhauser C, Negele RD, Ewringmann T (1991). Morphological effects of acute and chronic atrazine exposure in rainbow trout (*Oncorhynchus mykiss*). *Archives of Environmental Contamination and Toxicology*. 20, 454-461.
- Fontana L, Moreira E, Torres M, Fernandez I, Rios A, Sanchez DF, Gil A (1998). Dietary nucleotides correct plasma and liver microsomal fatty acids alterations in rats with liver cirrhosis induced by oral intake of thioacetamide. *J. Hepatol.* 28, 662-669.
- Fouda FM (2004). Hematological and biochemical studies on the effects of a biological and chemical pesticides on the Nile catfish, *Clarias gariepinus*. *J. Egyptian-German Soc. Zool.* 43, 77-97.
- Frigg M, Prabucki AL, Ruhdel EU (1990). Effect of dietary vitamin E levels on oxidative stability of trout filets. *Aquaculture*. 84, 145-158.
- Gad NS (2005). Impact of environmental pollution in the southern region of Lake Manzalah Egypt on some biochemical parameters of Tilapia Zillii. *J. Egyptian-German Soc. Zool.* 48, 279-298.
- Geret F, Jouan A, Turpin V, Bebianno MJ, Cosson RP (2002). Influence of metal exposure on metallothionein synthesis and lipid peroxidation in two bivalve mollusks: the oyster (*Crassostrea gigas*) and the mussel (*Mytilus edulis*). *Aquatic Living Resources*. 15, 16-66.
- Haberer K, Normann S (1988). Pflanzenschutzmittel aus der Sicht der of-fentlichen Wasserversorgung, Teil II. *Wasser Boden*. 4, 258-264.
- Halver JE (2002). The vitamins. In: J. E. Halver, R. W. Hardy, (Eds.), *Fish Nutrition*. Academic Press, San Diego, CA, pp. 61-141.
- Hamre K, Berge RK, Lie O (1998). Turnover of α -, β -, and δ -tocopherol and distribution in subcellular and lipoprotein fractions indicate presence of and hepatic tocopherol binding protein in Atlantic salmon (*Salmo salar* L.). *Fish Physiology and Biochemistry*. 18, 71-83.
- Heath AVG (1987). *Water pollution and fish physiology*. CRC press, Boca Raton, Florida, USA.
- Heber D, Lu QY (2002). Overview of mechanisms of action of lycopene. *Experimental Biology and Medicine*. 227, 920-923.
- Huang CH, Chang RJ, Huang SL, Chen W (2003). Dietary vitamin E supplementation affects tissue lipid peroxidation of hybrid tilapia, *Oreochromis niloticus* x *O. aureus*. *Comparative Biochemistry and Physiology. Part B, Biochemistry and Molecular Biology*. 134, 265-270.
- Huang SL, Weng YM, Huang CH (2004). Lipid peroxidation in sarcoplasmic reticulum and muscle of tilapia is inhibited by dietary vitamin E supplementation. *Journal of Food Biochemistry*. 28, 101-111.
- Hung SSO, Slinger SJ (1982). Effect of dietary vitamin E on rainbow trout (*Salmo gairdneri*) muscle alpha-tocopherol and storage stability. *Int. J. Vitamin and Nutr. Res.* 52, 119-123.
- Hussein SY, Mekkawy IAA (2001). The effect of lead-exposure and lead-clay interaction on the growth performance, biochemical and physiological characteristics and histopathology of Tilapia zillii. *Bull. Fac.Sci., Assiut Univ.* 30, 65-97.
- James R, Sampath K (1999). Effect of the ion-exchanging agent, Zeolite, on reduction of cadmium toxicity: an experimental study on growth and elemental uptake in *Heteropneustes fossilis* (Bloch). *J. Aquaculture in the Tropics*. 14, 65-74.
- Joseph B, Raj SJ (2011). Impact of Pesticide Toxicity on Selected Biomarkers in Fishes. *Int. J. Zool. Res.* 7, 212-222.
- Kalender S, Ogutcu A, Uzunhisarcikli M, Acikgoz F, Durak D, Ulusoy Y, Kalender Y (2005). Diazinon-induced hepatotoxicity and protective effect of vitamin E on some biochemical indices and ultrastructural changes. *Toxicology*. 211, 197-206.
- Keith FP, Weber JL (1979). The effect of carbon tetrachloride on the total plasma protein concentration of rainbow trout, *Salmo gairdneri*. *Comparative Biochemistry and Physiology*. 63, 37-42.
- Khadre SEM (1988). The effect of experimentally induced inflammation on the blood pattern and haemopoietic organs of the teleost, *Clarias lazera*. *Bulletin of the Institute of Oceanography and Fisheries*. 14, 191-203.
- Khidr BM, Wassif ET, Hussein SY, Mekkawy IAA (2001). Studies on the effect of the herbicide atrazine on the liver and kidney of the freshwater catfish *Chrysichthys auratus*. *J. Egypt Ger. Soc. Zool.* 34, 283-300.
- Köprücü SS, Köprücü K, Ural MS, İspir U, Pala M (2006). Acute toxicity of organophosphorous pesticide diazinon and its effects on behavior and some hematological parameters of fingerling European catfish (*Silurus glanis* L.). *Pesticide Biochemistry and Physiology*. 86, 99-105.
- Marie MAS, Mohamed HA, El-Badawy AA (1998). Physiological and biochemical responses of the common carp, *Cyprinus carpio*, to an organophosphorous insecticide "profenofos". *Egyptian J. Zool.* 31, 279-302.
- Martins DA, Afonso LOB, Hosoya S, Lewis-McCrea LMP, VLM, Lall SP (2007). Effects of moderately oxidized dietary lipid and the role of vitamin E on the stress response in Atlantic halibut (*Hippoglossus hippoglossus* L.). *Aquaculture*. 272, 573-580.
- Mazeaud MM, Mazeaud F, Donaldson EM (1977). Primary and secondary effects of stress in fish, some new data with a general review. *Transactions of the American Fisheries Society*. 106, 201-207.
- Mekkawy IAA, Mahmoud UM, Wassif ET, Naguib M (2012). Protective Roles of Tomato Paste and Vitamin E on Cadmium-induced Histological and Histochemical Changes of Liver of *Oreochromis niloticus* (Linnaeus, 1758). *Journal of Fisheries and Aquatic Science*. 7, 240-265.
- Mekkawy IAA, Hussein SY, Abd El-Nasser M, Ahmed SM (1996). Comparative studies on the effects of herbicide atrazine on some blood constituents and protein electrophoretic patterns of *Oreochromis niloticus* and *Chrysichthys auratus* at Assiut, Egypt. *J. Egyptian-German Society of Zoology*. 19, 283-319.
- Mekkawy IAA, Mahmoud UM, Wassif ET, Naguib M (2011). Effects of Cadmium on Some Hematological and Biochemical Characteristics of *Oreochromis niloticus* (Linnaeus, 1758) Dietary Supplemented with Tomato paste and vitamin E. *J. Fish Physiol. Biochem.*
- Montero D, Tort L, Robaina L, Vergara JM, Izquierdo MS (2001). Low vitamin E in diet reduces stress resistance of gilthead seabream (*Sparus aurata*) juveniles. *Fish and Shellfish Immunology*. 11, 473-490.
- Moreira EAM, Fagundes RLM, Filho DW, Neves D, Sell F, Bellisle F, Kupek E (2005). Effects of diet energy level and tomato powder consumption on antioxidant status in rats. *Clinical Nutrition*. 24, 1038-1046.
- Nwani CD, Lakra WS, Nagpure NS, Kumar R, Kushwaha B, Srivastava SK (2010). Toxicity of the Herbicide Atrazine: Effects on Lipid Peroxidation and Activities of Antioxidant Enzymes in the Freshwater Fish *Channa Punctatus* (Bloch). *Int. J. Environ. Res. Public Health*. 7.
- Oehmichen U, Haberer K (1986). Stickstoffherbizide im Rhein. *Vom Wasser*. 66, 225-241.
- Ogur R, Coskun O, Korkmaz A, Oter S, Yaren H, Hasde M (2005). High nitrate intake impairs liver functions and morphology in rats; protective effects of α -tocopherol. *Environmental Toxicology and Pharmacology*. 20, 161-166.
- Ohkawa H, Ohishi N, Nagi K (1979). Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Analytical Biochemistry*. 95, 351-358.
- Okajima E, Tsutsumi M, Ozono S, Akai H, Denda A, Nishino H, Oshima

- S, Sakamoto H, Konishi Y (1998). Inhibitory Effect of Tomato Juice on Rat Urinary Bladder Carcinogenesis after N-Butyl-N-(4-hydroxybutyl)nitrosamine Initiation. *Jpn. J. Cancer Res.* 89, 22-26.
- Okayi RG, Tachia MU, Ataguba GA, Dikwaha SH (2013a). Toxicity of the Herbicide Propanil on *Oreochromis niloticus* Fingerlings. *Journal of Fisheries and Aquatic Science* 8, 233-237.
- Okayi RG, Tachia MU, Ataguba GA, Dikwahal SH (2013b). Toxicity of the Herbicide Propanil on *Oreochromis niloticus* Fingerlings. *J. Fisheries and Aquatic Sci.* 8, 233-237.
- Ovie KS, Mohammed AK, Theresa MI (2007). Acute Haematological Effects of Sublethal Levels of Paraquat on the African Catfish, *Clarias gariepinus* (Osteichthyes: Clariidae). *Res. J. Environ. Sci.* 1, 331-335.
- Pickering AD (1981). Stress and compensation in teleostean fishes. Response to social and physical factors. In: A. D. Pickering, (Ed.), *Stress and Fish*. Academic Press, New York, London.
- Ranganna S (1976). *Manual of Analysis of Fruits and Vegetable Products*. McGraw Hill, New Delhi.
- Romeo M, Gnassia-Baralli M (1997). Effect of heavy metals on lipid peroxidation in the Mediterranean clam, *Ruditapes decussatus*. *Comparative Biochemistry and Physiology*. 118, 33-37.
- Romeo, M., Bennani, N., Gnassia-Barelli, M., Lafaurie, M., Girard, J. P., 2000. Cadmium and copper display different responses towards oxidative stress in the kidney of the sea bass *Dicentrarchus labrax*. *Aquatic Toxicology*. 48, 185-194.
- Saha S, Kaviraj A (2003). Acute toxicity synthetic pyrethroid cypermethrin freshwater catfish, *Heteropneustes fossilis* (Block). *Int. J. Toxicol.* 22, 325-328.
- Salgo MG, Corongiu FP, Sevnian A (1993). Enhanced interfacial catalysis and hydrolytic specificity of phospholipase A2 toward peroxidized phosphatidylcholine vesicles. *Archives of Biochemistry and Biophysics*. 304, 123-132.
- Sarkar S, Yadav P, Bhatnagar D (1997). Cadmium-induced lipid Peroxidation and the antioxidant system in rat erythrocytes: the role of antioxidants. *J. Trace Elements in Med. Biol.* 11, 8-13.
- Scaife JR, Onibi GE, Murray I, Fletcher TC, Houlihan DF (2000). Influence of α -tocopherol acetate on the short- and longterm storage properties of fillets from Atlantic salmon *Salmo salar* fed a high lipid diet. *Aquaculture Nutrition*. 6, 65-71.
- Shah SL (2006). Haematological parameters in tench, *Tinca tinca* after short term exposure to lead. *J. Appl. Toxicol.* 26, 223-228.
- Shalaby AME (2007). Effect of EDTA on reduction of cadmium toxicity on growth, some haematological and biochemical profiles of Nile tilapia (*Oreochromis niloticus*). *J. Fisheries Aquatic Sci.* 2, 100-109.
- Siang HY, Yee LM, Seng CT (2007). Acute toxicity of organochlorine insecticide endosulfan and its effect on behaviour and some haematological parameters of Asian swamp eel (*Monopterus albus*, Zuiew). *Pesticide Biochemistry and Physiology*. 89, 46-53.
- Sobh K, Poornima A, Harini P, Veeraiyah K (2007). A study on biochemical changes in the fresh water fish, *Catla catla* (Hamilton) exposed to the heavy metals toxicant cadmium chloride. *Kathmandu University. Journal of Science, Engineering and Technology*. 1.
- SPSS (1998). *SPSS for Windows*. Release 9.0.0 SPSS Inc.
- Stahl, W., Sies, H., 2003. Antioxidant activity of carotenoids. *Molecular Aspects of Medicine*. 24, 345-351.
- Stoskopf MK (1993). *Fish Medicine*. W. B. Saunders Company Harcour Brace Jovanovich, Inc.
- Tapiero H, Townsend DM, Tew KD (2004). The role of carotenoids in the prevention of human pathologies. *Biomedicine and Pharmacotherapy*. 58, 100-110.
- Tasli S, Patty L, Boetti H, Ravanel P, Vachaud G, Scharif C, al., e (1996). Persistence and leaching of atrazine in corn culture in the experimental site of La Cote Saint Andre´ (Ise`re, France). *Archives of Environment Contamination and Toxicology*. 30, 203-212.
- Ursini F, Maiorino M, Sevanian A (1991). Membrane hydroperoxides in oxidative stress. In: H. Sies, (Ed.), *Oxidants and Antioxidants*. Harcourt Brace Jovanovich, London, pp. 319-336.
- Vankampen EJ (1961). Determination of hemoglobin. *Clinica Chimica Acta*. 6, 538-544.
- Velmurugan B, Bhuvanewari V, Balasenthil S, Nagini S (2001). Lycopene, an antioxidant carotenoid modulates glutathione-dependent hepatic biotransformation enzymes during experimental gastric carcinogenesis. *Nutrition Research*. 21, 1117-1124.
- Velmurugan B, Bhuvanewari V, Nagini S (2002). Antiperoxidative effects of lycopene during N-methyl-N-nitro-nitrosoguanidine-induced gastric carcinogenesis. *Fitoterapia*. 73, 604-611.
- Viran R, Krkoç FU, Polat H, Koçak O (2003). Investigation of acute toxicity of deltamethrin on guppies (*Poecilia reticulata*). *Ecotoxicology and Environmental Safety*. 55, 82-85.
- Visioli F, Riso P, Grande S, Galli C, Porrini M (2003). Protective activity of tomato products on in vivo markers of lipid oxidation. *Eur. J. Nutr.* 42, 201-206.
- Wang Y, Yuen KH, Ng WK (2006). Deposition of tocotrienols and tocopherols in the tissues of red hybrid tilapia, *Oreochromis sp.*, fed γ -tocotrienol-rich fraction extracted from crude palm oil and its effect on lipid peroxidation. *Aquaculture*. 253, 583-591.
- Wassif ET, Khidr BM, Hussein SY, Mekkawy IAA, Hassan HI (2000). Effects of herbicide atrazine on the structure of some organs of the Nile fish *Oreochromis niloticus*. The 4th Conference of the Egyptian Society for the Development of Fisheries Resources and Human Health, Ain Shams University, Faculty of Science, Cairo.
- Wertz K, Siler U, Goralczyk R (2004). Lycopene: modes of action to promote prostate health. *Archives of Biochemistry and Biophysics*. 430, 127-134.
- WHO/FAO (1996). *WHO/FAO data sheets on pesticides (Atrazine)*. WHO/PCS/DS/96. 82, 1-18.
- Wintrobe MM (1978). *Clinical hematology*. Henry Kimpton, London, 448.
- Yadav AS, Bhatnagar A, Kaur M (2010). Assessment of Genotoxic Effects of Butachlor in Fresh Water Fish, *Cirrhinus mrigala* (Hamilton). *Research Journal of Environmental Toxicology*. 4, 223-230.
- Yilmaz O, Celik S, Cay M, Naziroglu M (1997). Protective Role of Intraperitoneally Administrated Vitamin E and Selenium on the Levels of Total Lipid, Total Cholesterol and Fatty Acid Composition of Muscle and Liver Tissues in Rats. *J. Cellular Biochem.* 64, 233-241.
- Yilmaz S, Atessahin A, Sahna E, Karahan I, Ozer S (2006). Protective effect of lycopene on adriamycin-induced cardiotoxicity and nephrotoxicity. *Toxicology*. 218, 164-171.